

Product datasheet for AP14723PU-N

ALDH4A1 (C-term) Rabbit Polyclonal Antibody

Product data:

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product Type:	Primary Antibodies
Applications:	IHC, WB
Recommended Dilution:	ELISA: 1/1,000. Western blotting: 1/50 - 1/100. Immunohistochemistry: 1/50 - 1/100.
Reactivity:	Human
Host:	Rabbit
lsotype:	lg
Clonality:	Polyclonal
Immunogen:	This antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide selected from the C-terminal region of human ALDH4A1.
Specificity:	This antibody reacts to ALDH4A1.
Formulation:	PBS with 0.09% (W/V) sodium azide State: Purified State: Liquid purified Ig
Concentration:	lot specific
Purification:	Prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS
Conjugation:	Unconjugated
Storage:	Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Gene Name:	aldehyde dehydrogenase 4 family member A1
Database Link:	<u>Entrez Gene 8659 Human</u> <u>P30038</u>




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

GRIGENE ALDH4A1 (C-term) Rabbit Polyclonal Antibody – AP14723PU-N


Background:ALDH4A1 belongs to the aldehyde dehydrogenase family of proteins. This enzyme is a
mitochondrial matrix NAD-dependent dehydrogenase which catalyzes the second step of the
proline degradation pathway, converting pyrroline-5-carboxylate to glutamate. Deficiency of
this enzyme is associated with type II hyperprolinemia, an autosomal recessive disorder
characterized by accumulation of delta-1-pyrroline-5-carboxylate (P5C) and proline.

Synonyms: ALDH4A1, P5CDH, P5C dehydrogenase

Product images:



Western blot analysis of anti-ALDH4A1 Antibody (C-term) in HepG2 cell line lysates (35ug/lane). ALDH4A1 (arrow) was detected using the purified Pab.



Formalin-fixed and paraffin-embedded human hepatocarcinoma tissue reacted with ALDH4A1 antibody (C-term), which was peroxidaseconjugated to the secondary antibody, followed by DAB staining.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US