

## **Product datasheet for AP14979PU-N**

## 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

CN: techsupport@origene.cn

OriGene Technologies, Inc.

## **BRD3 (C-term) Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type:** Primary Antibodies

**Applications:** IHC, WB

Recommended Dilution: ELISA: 1/1,000.

Western blotting: 1/100 - 1/500.

Immunohistochemistry: 1/50 - 1/100.

Reactivity: Human, Mouse

**Host:** Rabbit

**Isotype:** lg

Clonality: Polyclonal

**Immunogen:** This antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide

selected from the C-terminal region of human BRD3.

**Specificity:** This antibody reacts to BRD3.

**Formulation:** PBS with 0.09% (W/V) sodium azide

State: Purified

State: Liquid purified Ig

**Concentration:** lot specific

**Purification:** Protein G column, eluted with high and low pH buffers and neutralized immediately, followed

by dialysis against PBS

Conjugation: Unconjugated

Storage: Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

**Stability:** Shelf life: one year from despatch.

Gene Name: bromodomain containing 3

Database Link: Entrez Gene 8019 Human

Q15059

**Background:** This gene was identified based on its homology to the gene encoding the RING3 protein, a

serine/threonine kinase. The gene localizes to 9q34, a region which contains several major histocompatibility complex (MHC) genes. The function of the encoded protein is not known.







Synonyms:

KIAA0043, RING3L, RING3-like protein, Bromodomain-containing protein 3

## **Product images:**



The anti-BRD3 Pab is used in Western blot to detect BRD3 in mouse brain tissue lysate (Lane 1) and Saos-2 cell lysate (Lane 2).



Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by AEC staining.