

# Product datasheet for AP15225PU-N

### PTP epsilon (PTPRE) (C-term) Rabbit Polyclonal Antibody

### **Product data:**

#### **Product Type: Primary Antibodies** WB **Applications:** Recommended Dilution: ELISA: 1/1,000. Western blotting: 1/100 - 1/500. **Reactivity:** Human Host: Rabbit Isotype: lg **Clonality:** Polyclonal This antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide Immunogen: selected from the C-terminal region of human PTPepsilon. Specificity: This antibody reacts to PTPepsilon. Formulation: PBS with 0.09% (W/V) sodium azide State: Purified State: Liquid purified Ig **Concentration:** lot specific **Purification:** Protein G column, eluted with high and low pH buffers and neutralized immediately, followed by dialysis against PBS **Conjugation:** Unconjugated Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Storage: Avoid repeated freezing and thawing. Stability: Shelf life: one year from despatch. Gene Name: protein tyrosine phosphatase, receptor type E Database Link: Entrez Gene 5791 Human P23469



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

	PTP epsilon (PTPRE) (C-term) Rabbit Polyclonal Antibody – AP15225PU-N
Background:	PTPepsilon is a member of the protein tyrosine phosphatase (PTP) family. PTPs are known to be signaling molecules that regulate a variety of cellular processes including cell growth, differentiation, mitotic cycle, and oncogenic transformation. Two alternatively spliced transcript variants of the gene have been reported, one of which encodes a receptor-type PTP that possesses a short extracellular domain, a single transmembrane region, and two tandem intracytoplasmic catalytic domains; Another one encodes a PTP that contains a distinct hydrophilic N-terminus, and thus represents a nonreceptor-type isoform of this PTP. Studies of the similar gene in mice suggested the regulatory roles of this PTP in RAS related signal transduction pathways, cytokines induced SATA signaling, as well as the activation of voltage- gated K+ channels.

**Synonyms:** Protein-tyrosine phosphatase epsilon, R-PTP-epsilon

## Product images:



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US