

Product datasheet for **AP17780PU-N**

Thymine DNA glycosylase (TDG) (C-term) Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	WB
Recommended Dilution:	ELISA: 1/1,000. Western blotting: 1/50 - 1/100.
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	KLH conjugated synthetic peptide selected from the C-terminal region of human TDG
Specificity:	This antibody reacts to TDG.
Formulation:	PBS with 0.09% (W/V) sodium azide State: Liquid purified Ig
Concentration:	lot specific
Purification:	Saturated Ammonium Sulfate (SAS) precipitation
Conjugation:	Unconjugated
Storage:	Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Gene Name:	thymine DNA glycosylase
Database Link:	Entrez Gene 6996 Human Q13569
Background:	The protein TDG belongs to the TDG/mug DNA glycosylase family. Thymine-DNA glycosylase (TDG) removes thymine moieties from G/T mismatches by hydrolyzing the carbon-nitrogen bond between the sugar-phosphate backbone of DNA and the mispaired thymine. With lower activity, this enzyme also removes thymine from C/T and T/T mispairings. TDG can also remove uracil and 5-bromouracil from mispairings with guanine. This enzyme plays a central role in cellular defense against genetic mutation caused by the spontaneous deamination of 5-methylcytosine and cytosine.
Synonyms:	E130317C12Rik; hTDG; JZA-3; Jza1; OTTMUSP00000028912; OTTMUSP00000028913




[View online »](#)

Protein Families: Druggable Genome

Protein Pathways: Base excision repair

Product images:



Western blot analysis of TDG antibody (C-term) in K562 cell line lysates (35ug/lane). TDG (arrow) was detected using the purified Pab.