

## Product datasheet for AP20194PU-M

## 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

CN: techsupport@origene.cn

OriGene Technologies, Inc.

## **COX2 (PTGS2) Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type:** Primary Antibodies

Applications: IF, IHC, WB

Recommended Dilution: Western blot: 1/500-1/1000.

Immunohistochemistry on paraffin sections: 1/50-1/200.

Immunofluorescence: 1/50-1/200.

Reactivity: Human
Host: Rabbit

Clonality: Polyclonal

**Specificity:** This antibody antibody detects endogenous levels of Cox2/PGHS2 protein.

**Formulation:** Phosphate buffered saline (PBS), pH~7.2 containing 0.05% Sodium Azide as preservative.

State: Aff - Purified

State: Liquid purified Ig fraction (> 95% pure by SDS-PAGE).

Concentration: 1.0 mg/ml

**Purification:** Affinity Chromatography using epitope-specific immunogen.

Conjugation: Unconjugated

**Storage:** Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

**Stability:** Shelf life: one year from despatch.

Predicted Protein Size: ~ 74 kDa

**Gene Name:** prostaglandin-endoperoxide synthase 2

**Database Link:** Entrez Gene 5743 Human

P35354



Background:

Prostaglandins are a diverse group of autocrine and paracrine hormones that mediate many cellular and physiologic processes. Prostaglandin H2 (PGH2) is an intermediate in formation of the prostaglandins. Two prostaglandin synthases that catalyze the formation of PGH2 from arachidonic acid (AA) are cyclooxygenase-1 and cyclooxygenase-2. Cyclooxygenase-2, or Cox-2, is efficiently induced in migratory cells responding to pro-inflammatory stimuli and is considered to be an important mediator of inflammation. An alternative form of the protein, designated Cox-1, is constitutively expressed in most tissues and is thought to serve in general "housekeeping" functions. Both enzymes are targets for the nonsteroidal therapeutic anti-inflammatorydrugs, NSAIDs.


Synonyms: Cyclooxygenase-2, PTGS2, COX2, PHS II, PGH synthase 2, PGHS-2, PGHS2

**Protein Families:** Druggable Genome

**Protein Pathways:** Arachidonic acid metabolism, Pathways in cancer, Small cell lung cancer, VEGF signaling

pathway


## **Product images:**



Western blot (WB) analysis of Cox2/PGHS2 (pThr594) antibody (Cat.-No.: [AP20194PU-N])

Lane1: Hela whole cell lysate treated with Lps
Lane2: Jurkat whole cell lysate treated with Lps
Lane3: Raw264.7 whole cell lysate treated with Lps
Lane4: Rat lung tissue lysate

Lane4: Rat lung tissue lysate Cox-2 (T594) pAb at 1:500 dilution



Immunohistochemistry (IHC) analyzes of Cox2/PGHS2 (pThr594) antibody (Cat.-No.: [AP20194PU-N]) in paraffin-embedded human breast carcinoma tissue.