

Product datasheet for AP20207PU-N

B MyB (MYBL2) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IF, WB

Recommended Dilution: Western Blot: 1/500-1/1000.

Immunofluorescence: 1/50-1/200.

Reactivity: Human, Mouse, Rat

Host: Rabbit

Clonality: Polyclonal

Specificity: This antibody antibody detects endogenous levels of B-Myb / MYB2 protein.

Formulation: Phosphate buffered saline (PBS), pH~7.2

State: Aff - Purified

State: Liquid purified Ig fraction (> 95% pure by SDS-PAGE)

Preservative: 0.05% Sodium Azide

Concentration: 1.0 mg/ml

Purification: Affinity Chromatography using epitope-specific immunogen

Conjugation: Unconjugated

Storage: Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: ~80 kDa

Gene Name: MYB proto-oncogene like 2

Database Link: Entrez Gene 17865 MouseEntrez Gene 296344 RatEntrez Gene 4605 Human

P10244



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Background:

Expression of the B-Myb transcription factor is upregulated during late G1 phase of the cell cycle by an E2F-dependent transcriptional mechanism. B-Myb is specifically phosphorylated during S phase, suggesting that a cyclin-dependent kinase (Cdk) regulates its activity. Consistent with this notion, the S phase-specific cyclin A/Cdk2 was found previously to enhance B-Myb transactivation activity in cotransfected cells. There is evidence that B-Myb is a direct physiological target for cyclin A/Cdk2. Data indicate that phosphorylation by cyclin A/Cdk2 is directly involved in enhancing B-Myb transactivation activity and that levels of endogenous cyclin A/Cdk2 activity may contribute to cell line-specific B-Myb function . Nuclear entry of B-Myb is dependent on multiple nuclear localization signals (NLS's). Mutagenesis of the putative NLS's of B-Myb has identified two separate NLS's, NLS1 and NLS2. Each of the two NLS's is essential for efficient nuclear targeting.

Synonyms: BMYB, B-Myb

Protein Families: Druggable Genome, Stem cell - Pluripotency, Transcription Factors

Product images:



Western blot analysis of B-Myb antibody (Cat.-No. AP20207PU-N) in extracts from K562 cells.