

## **Product datasheet for AP20216PU-N**

## 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

OriGene Technologies, Inc.

## **CREB1 Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type:** Primary Antibodies

Applications: IHC, WB

Recommended Dilution: Western blot: 1/500-1/1000.

Immunohistochemistry on paraffin sections: 1/50-1/200.

Reactivity: Human, Mouse, Rat

**Host:** Rabbit

Clonality: Polyclonal

**Specificity:** This antibody detects endogenous levels of CREB protein.

**Formulation:** Phosphate buffered saline (PBS) with 0.05% sodium azide, approx. pH 7.2

State: Aff - Purified

State: Liquid purified Ig fraction

**Concentration:** 1.0 mg/ml

**Purification:** Affinity-chromatography using epitope-specific immunogen; purity is > 95% (by SDS-PAGE)

Conjugation: Unconjugated

**Storage:** Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

**Stability:** Shelf life: one year from despatch.

Predicted Protein Size: ~ 38 kDa

**Gene Name:** cAMP responsive element binding protein 1

Database Link: Entrez Gene 12912 MouseEntrez Gene 81646 RatEntrez Gene 1385 Human

P16220





Background:


Eukaryotic gene transcription is regulated by sequence-specific transcription factors that bind modular cis acting promoter and enhancer elements. The ATF/CREB transcription factor family binds the palindromic cAMP response element (CRE) octanucleotide TGACGTCA. The ATF/CREB family includes CREB-1, CREB-2 (also designated ATF-4), ATF-1, ATF-2 and ATF-3. This family of proteins contains highly divergent N-terminal domains, but shares a C-terminal leucine zipper for dimerization and DNA binding. Although CREB can bind to DNA in an unphosphorylated state, it cannot activate transcription. Phosphorylation of CREB on Ser 133 by protein kinase A facilitates its interaction with the CREB-binding protein (CBP) and activates the basal transcription complex. CREB functions in neoglucogenesis through interactions with the nuclear coactivator PGC-1. CREB may play a role in the pathogenesis of type II diabetes and dilated cardiomyopathy.

Synonyms: CREB-1


**Protein Families:** Druggable Genome, Transcription Factors

**Protein Pathways:** Antigen processing and presentation, Huntington's disease, Melanogenesis, Prostate cancer

## **Product images:**



Western blot analysis of CREB antibody (AP20216PU-N) in extracts from HeLa cells.



Immunohistochemistry analyzes of CREB antibody (AP20216PU-N) in paraffin-embedded human tonsil tissue.