

Product datasheet for AP20230PU-N

GATA2 (+GATA3) Rabbit Polyclonal Antibody

Product data:

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product Type:	Primary Antibodies
Applications:	IF, IHC
Recommended Dilution:	Immunofluorescence: 1/50-1/200. Immunohistochemistry on Paraffin Sections: 1/50-1/200.
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Specificity:	This antibody detects endogenous levels of GATA2/3 protein.
Formulation:	Phosphate buffered saline (PBS), pH~7.2 State: Aff - Purified State: Liquid purified Ig fraction (> 95% pure by SDS-PAGE Preservative: 0.05% Sodium Azide
Concentration:	1.0 mg/ml
Purification:	Affinity Chromatography using epitope-specific immunogen
Conjugation:	Unconjugated
Storage:	Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Gene Name:	GATA binding protein 2
Database Link:	<u>Entrez Gene 2624 Human</u> <u>P23769</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US


GATA2 (+GATA3) Rabbit Polyclonal Antibody – AP20230PU-N

Background:The genes for all 4 subunits of the T-cell antigen receptor (alph,a bet,a gamma and delta) are
controlled by distinct enhancers and their enhancer-binding proteins. Marine and Winoto
(1991) identified a common TCR regulatory element by demonstrating binding of the
enhancer-binding protein GATA3 to the enhancer elements of all 4 TCR genes. GATA3 had
been shown in the chicken to be an enhancer-binding protein containing a zinc finger
domain. GATA3 mRNA was demonstrated by Northern blot analysis in T cells but not in B
cells or macrophages. GATA3 is abundantly expressed in the T-lymphocyte lineage and is
thought to participate in T-cell receptor gene activation through binding to enhancers.
Labastie et al. (1994) cloned the human gene and the 5-prime end of the mouse gene. The
human gene comprises 6 exons distributed over 17 kb of DNA. Its 2 zinc fingers are encoded
by 2 separate exons highly conserved with those of GATA1, but no other structural
homologies between the 2 genes could be found.

Synonyms:

GATA-2, GATA-binding protein 2

Product images:



Immunohistochemistry analysis of GATA2/3 Antibody Cat.-No AP20230PU-N in paraffinembedded human placenta tissue.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US