

## Product datasheet for **AP20297PU-N**

### Alpha B Crystallin (CRYAB) Rabbit Polyclonal Antibody

#### Product data:

Product Type:	Primary Antibodies
Applications:	IHC
Recommended Dilution:	<b>Immunohistochemistry on paraffin sections:</b> 1/50-1/200.
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Specificity:	This antibody detects endogenous levels of Crystallin-αB protein.
Formulation:	Phosphate buffered saline (PBS) with 0.05% sodium azide, approx. pH 7.2 State: Aff - Purified State: Liquid purified Ig fraction
Concentration:	1.0 mg/ml
Purification:	Affinity-chromatography using epitope-specific immunogen; purity is > 95% (by SDS-PAGE)
Conjugation:	Unconjugated
Storage:	Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	~ 26 kDa
Gene Name:	crystallin alpha B
Database Link:	<a href="#">Entrez Gene 12955 Mouse</a> <a href="#">Entrez Gene 25420 Rat</a> <a href="#">Entrez Gene 1410 Human</a> <a href="#">P02511</a>




[View online »](#)

**Background:**

The baculovirus protein p35 inhibits virally-induced apoptosis of invertebrate and mammalian cells and may function to impair the clearing of virally infected cells by the immune system of the host. This is accomplished at least in part by the ability of p35 to block both TNF- and FAS-mediated apoptosis through the inhibition of the ICE family of serine proteases. Two mammalian homologs of baculovirus p35, referred to as inhibitor of apoptosis protein (IAP) 1 and 2, share an amino-terminal baculovirus IAP repeat (BIR) motif and a carboxy-terminal RING finger. Although the c-IAPs do not directly associate with the TNF receptor (TNF-R), they efficiently block TNF-mediated apoptosis through their interaction with the downstream TNF-R effectors, TRAF1 and TRAF2. Additional IAP family members include ILP (for IAP-like protein) and survivin. ILP inhibits activated caspase-3, leading to the resistance of FAS-mediated apoptosis. Survivin (also designated TIAP) is expressed during the G2/M phase of the cell cycle and associates with microtubules of the mitotic spindle. Increased caspase-3 activity is detected when a disruption of survivin-microtubule interactions occurs.

**Synonyms:**

Alpha(B)-crystallin, CRYAB, Heat shock protein beta-5, Renal carcinoma antigen NY-REN-27, HspB5

**Product images:**


Immunohistochemistry analyzes of Crystallin-αB antibody (AP20297PU-N) in paraffin-embedded human heart tissue.