

Product datasheet for **AP20307PU-M**

IKK alpha (CHUK) Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	IHC, WB
Recommended Dilution:	Western blot: 1/500-1/1000. Immunohistochemistry on paraffin sections 1/50-1/200.
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Specificity:	This antibody detects endogenous levels of IKK α /β protein.
Formulation:	Phosphate buffered saline (PBS) with 0.05% sodium azide, approx. pH 7.2 State: Aff - Purified State: Liquid purified Ig fraction
Concentration:	1.0 mg/ml
Purification:	Affinity-chromatography using epitope-specific immunogen; purity is > 95% (by SDS-PAGE)
Conjugation:	Unconjugated
Storage:	Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	~ 85 kDa
Gene Name:	conserved helix-loop-helix ubiquitous kinase
Database Link:	Entrez Gene 12675 Mouse Entrez Gene 309361 Rat Entrez Gene 1147 Human O15111



[View online »](#)


Background: The transcription factor NFκB is retained in the cytoplasm in an inactive form by the inhibitory protein IκB. Activation of NFκB requires that IκB be phosphorylated on specific serine residues, which results in targeted degradation of IκB. IκB kinase α (IKKα), previously designated CHUK, interacts with IκBα and specifically phosphorylates IκBα on Serine 32 and 36, the sites that trigger its degradation. IKKα appears to be critical for NFκB activation in response to proinflammatory cytokines. Phosphorylation of IκB by IKKα is stimulated by the NFκB inducing kinase (NIK), which itself is a central regulator for NFκB activation in response to TNF and IL-1. The functional IKK complex contains three subunits, IKKα, IKKβ and IKKγ (also designated NEMO), and each appear to make essential contributions to IκB phosphorylation.

Synonyms: CHUK, TCF16, I kappa-B kinase alpha, IκBKA, IKK-alpha, IKK-A, IκappaB kinase, I-kappa-B kinase 1, NFKB1A, IKK1


Protein Families: Druggable Genome, Protein Kinase

Protein Pathways: Acute myeloid leukemia, Adipocytokine signaling pathway, Apoptosis, B cell receptor signaling pathway, Chemokine signaling pathway, Chronic myeloid leukemia, Cytosolic DNA-sensing pathway, Epithelial cell signaling in Helicobacter pylori infection, MAPK signaling pathway, NOD-like receptor signaling pathway, Pancreatic cancer, Pathways in cancer, Prostate cancer, RIG-I-like receptor signaling pathway, Small cell lung cancer, T cell receptor signaling pathway, Toll-like receptor signaling pathway

Product images:



Western blot (WB) analysis of IKKα/β antibody (Cat.-No.: [AP20307PU-N]) at 1/500 dilution Lane 1:THP-1 whole cell lysateLane 2:HEK293T whole cell lysateLane 3:Mouse brain tissue lysateLane 4:Rat heart tissue lysateLane 5:NIH-3T3 whole cell lysate



Immunohistochemistry analyzes of IKK α / β antibody ([AP20307PU-N]) in paraffin-embedded human breast carcinoma tissue.