

## **Product datasheet for AP20324PU-N**

## 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

CN: techsupport@origene.cn

OriGene Technologies, Inc.

## **PAK1 Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type:** Primary Antibodies

Applications: IHC

Recommended Dilution: Immunohistochemistry on paraffin sections: 1/50-1/200.

Reactivity: Human, Mouse, Rat

Host: Rabbit
Clonality: Polyclonal

**Specificity:** This detects endogenous levels of PAK1/2/3 protein.

**Formulation:** Phosphate buffered saline (PBS) with 0.05% sodium azide, approx. pH 7.2

State: Aff - Purified

State: Liquid purified Ig fraction

Concentration: 1.0 mg/ml

**Purification:** Affinity-chromatography using epitope-specific immunogen; purity is > 95% (by SDS-PAGE)

**Conjugation:** Unconjugated

Storage: Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

**Stability:** Shelf life: one year from despatch.

**Predicted Protein Size:** ~ 60 kDa

Gene Name: p21 (RAC1) activated kinase 1

Database Link: Entrez Gene 18479 MouseEntrez Gene 29431 RatEntrez Gene 5058 Human

Q13153



Background:

Three isoforms of serine/threonine kinases, designated  $\alpha$ PAK p68,  $\beta$ PAK p65 and  $\gamma$ PAK p62, have been shown to exhibit a high degree of sequence homology with the S. cerevisiae kinase Ste 20, involved in pheromone signaling. The  $\alpha$ ,  $\beta$  and  $\gamma$ PAK isoforms complex specifically with Rac1 and Cdc42 in their active GTP-bound state, inhibiting their intrinsic GTPase activity leading to their autophosphorylation. There are eight sites of autophosphorylation on  $\gamma$ PAK, including Ser 19, Ser 141 and Thr 402, and phosphorylation of Ser 141 and Thr 402 is correlated with  $\gamma$ PAK activation. Once phosphorylated and their affinity for Rac/Cdc42 reduced, the PAK isoforms disassociate from the complex to seek downstream substrates. One such putative substrate is Mek kinase, an upstream effector of Mek4 which is involved in the JNK signaling pathway. While the PAK isoforms interact in a GTPdependent manner with Rac1 and Cdc42, they do not interact with Rho.

Synonyms:

PAK 1, PAK-1, Alpha-PAK, PAK alpha, p21-activated kinase 1, p65-PAK, PAK 2, PAK-2, Gamma-PAK, PAK gamma, PAK65, p21-activated kinase 2, p58, PAK 3, PAK-3, Beta-PAK, PAK beta, p21-activated kinase 3, Oligophrenin-3, OPHN3


**Protein Families:** 

Druggable Genome, Protein Kinase, Stem cell - Pluripotency

**Protein Pathways:** 

Axon guidance, Chemokine signaling pathway, Epithelial cell signaling in Helicobacter pylori infection, ErbB signaling pathway, Fc gamma R-mediated phagocytosis, Focal adhesion, MAPK signaling pathway, Natural killer cell mediated cytotoxicity, Regulation of actin cytoskeleton, Renal cell carcinoma, T cell receptor signaling pathway

## **Product images:**



Immunohistochemistry analyzes of PAK1/2/3 antibody (AP20324PU-N) in paraffin-embedded human brain tissue.