

Product datasheet for AP20345PU-M

Delta Opioid Receptor (OPRD1) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies IHC, WB **Applications: Recommended Dilution: Western blot:** 1/500-1/1000. Immunohistochemistry on Paraffin Sections: 1/50-1/200. **Reactivity:** Human, Mouse, Rat Host: Rabbit **Clonality:** Polyclonal Specificity: This antibody detects endogenous levels of DOR-1 protein. Formulation: Phosphate buffered saline (PBS), pH~7.2 State: Aff - Purified State: Liquid purified Ig fraction (> 95% pure by SDS-PAGE) Preservative: 0.05% Sodium Azide **Concentration:** 1.0 mg/ml **Purification:** Affinity Chromatography using epitope-specific immunogen **Conjugation:** Unconjugated Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Storage: Avoid repeated freezing and thawing. Shelf life: one year from despatch. Stability: **Predicted Protein Size:** ~ 35, 50 kDa Gene Name: opioid receptor delta 1 Database Link: Entrez Gene 18386 MouseEntrez Gene 24613 RatEntrez Gene 4985 Human P41143

View online »

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

OriGene Technologies, Inc.


9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn



	Delta Opioid Receptor (OPRD1) Rabbit Polyclonal Antibody – AP20345PU-M
Background:	The opioid receptors are G-protein coupled, seven-transmembrane domain receptors for enkephalins, endorphins, and dynorphins. Three different opioid receptor subtypes (kappa, delta and mu) were first identified by their different selectivities for various naturally occurring alkaloid opioid ligands, and subsequently confirmed by molecular cloning. The amino acid sequences of the opioid receptor subtypes are ~70% homologous, and are similar to somatostatin receptors (SSTRs) showing ~40 % homology with SSTR1. G-protein binding is thought to occur at the third intracellular loop of the opioid receptors, which is also the location of consensus sequences for phosphorylation of the receptor. Interestingly, the genes encoding the specific receptor subtypes are found on different chromosomes in both the human and mouse genomes.
Synonyms:	OPRD1, OPRD, DOR-1, D-OR-1
Protein Families:	Druggable Genome, Transmembrane
Protein Pathways	Neuroactive ligand-receptor interaction

Product images:

~ \$



Western blot analysis of DOR-1 antibody (Cat.-No [AP20345PU-N]) in extracts from HUVEC cells.



Immunohistochemistry analysis of DOR-1 antibody (Cat.-No [AP20345PU-N]) in paraffinembedded human brain tissue.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US