

Product datasheet for AP20371PU-N

TRF1 (TERF1) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: Western blot: 1/500-1/1000.

Immunohistochemistry on paraffin sections: 1/50-1/200.

Reactivity: Human, Mouse

Host: Rabbit

Clonality: Polyclonal

Specificity: This antibody detects endogenous levels of TRF1 protein.

Formulation: Phosphate buffered saline (PBS) with 0.05% sodium azide, approx. pH 7.2

State: Aff - Purified

State: Liquid purified Ig fraction

Concentration: 1.0 mg/ml

Purification: Affinity-chromatography using epitope-specific immunogen; purity is > 95% (by SDS-PAGE)

Conjugation: Unconjugated

Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Storage:

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: ~ 50 kDa

Gene Name: telomeric repeat binding factor 1

Database Link: Entrez Gene 21749 MouseEntrez Gene 7013 Human

P54274



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Background:

Telomeric repeat binding factor 1 (TRF1, TERF1, PIN2, TRBF1) and telomeric repeat binding factor 2 (TRF2, TERF2, TRBF2) are present at telomeres throughout the cell cycle, where they regulate telomerase by acting in cis to limit the elongation of individual chromosome ends. Telomerase adds hexameric repeats of 5'-TTAGGG-3' to the ends of chromosomal DNA. This telomerase enzyme plays an influential role in cellular immortalization and cellular senescence. TRF1 negatively regulates telomere elongation, while TRF2 protects the chromosome ends by inhibiting end-to-end fusions. Downregulation of TRF expression in tumor cells may contribute to cell immortalization and malignant progression. TRF1 has an acidic N-terminus while TRF2 has a basic N-terminus. TRF2 localizes in the nucleolus at G0 and S and diffuses out of the nucleolus in G2 phase. During mitosis TRF2 disperses from the condensed chromosomes and returns to the nucleolus at cytokinesis.


Synonyms: PIN2, TRBF1, TRF, TRF1

Protein Families: Transcription Factors

Product images:



Western blot analysis of TRF1 antibody (AP20371PU-N) in extracts from 293 cells.



Immunohistochemistry analyzes of TRF1 antibody (AP20371PU-N) in paraffin-embedded human brain tissue.