

## Product datasheet for AP20384PU-N

## **STAT1 Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type: Primary Antibodies** 

**Applications:** IHC, WB

Recommended Dilution: Western Blot: 1/500-1/1000.

Immunohistochemistry on paraffin sections: 1/50-1/200.

Reactivity: Human Host: Rabbit

Clonality: Polyclonal

Specificity: This antibody detects endogenous levels of Stat1 protein.

(region surrounding Leu177)

Formulation: Phosphate buffered saline (PBS) with 15 mM sodium azide, approx. pH 7.2.

State: Aff - Purified

State: Liquid purified Ig fraction

Concentration: 1.0 mg/ml

**Purification:** Affinity chromatography (> 95% (by SDS-PAGE)

Conjugation: Unconjugated

Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Storage:

Avoid repeated freezing and thawing.

Shelf life: one year from despatch. Stability:

~ 87 kDa **Predicted Protein Size:** 

Gene Name: signal transducer and activator of transcription 1

**Database Link:** Entrez Gene 6772 Human

P42224



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Background:

STATs (signal transducers and activators of transcription) were originally discovered as two proteins (STAT1 and STAT2) which were involved in interferon alpha (IFN alpha) and IFN gamma signal transduction. Since then, several additional STAT proteins have been identified (STAT3, 4, 5,a 5b, and 6). STATs undergo tyrosine phosphorylation in response to growth factor or cytokine signaling. This phosphorylation results in dimerization and translocation of STAT proteins to the nucleus. In some cases this process is mediated by JAK Kinases (Janus Kinases 1, 2, and 3). For maximum activation of these proteins, phosphorylation at specific tyrosine and serine residues may be required in STAT1 alph,a 3, 4, and 5. Specific functions of the STAT proteins are poorly defined at this time. They are thought to be involved in a variety of systems including antiviral responses and cell transformation.


**Synonyms:** DKFZp686B04100; ISGF-3; OTTHUMP00000205845; STAT91

**Protein Families:** Druggable Genome, Transcription Factors

**Protein Pathways:** Chemokine signaling pathway, Jak-STAT signaling pathway, Pancreatic cancer, Pathways in

cancer, Toll-like receptor signaling pathway

## **Product images:**



Hela whole cell lysate Stat1(L177) pAb at 1:500 dilution Western blot (WB) analysis of Stat1 antibody (Cat.-No.: AP20384PU-N) in extracts from hela cells.