

## Product datasheet for **AP20411PU-M**

### PFKFB1 (Liver) Rabbit Polyclonal Antibody

#### Product data:

Product Type:	Primary Antibodies
Applications:	IHC, WB
Recommended Dilution:	<b>Western blot:</b> 1/500-1/1000. <b>Immunohistochemistry on Paraffin Sections:</b> 1/50-1/200.
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Specificity:	This antibody detects endogenous levels of PFK-2 liv protein. (region surrounding Glu442)
Formulation:	Phosphate buffered saline (PBS), pH~7.2 containing 15 mM Sodium Azide as preservative. State: Aff - Purified State: Liquid purified Ig fraction (> 95% pure by SDS-PAGE).
Concentration:	1.0 mg/ml
Purification:	Affinity-Chromatography using epitope-specific immunogen.
Conjugation:	Unconjugated
Storage:	Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	~ 52 kDa
Gene Name:	6-phosphofructo-2-kinase/fructose-2,6-biphosphatase 1
Database Link:	<a href="#">Entrez Gene 18639 Mouse</a> <a href="#">Entrez Gene 24638 Rat</a> <a href="#">Entrez Gene 5207 Human</a> <a href="#">P16118</a>



[View online »](#)

**Background:**

Phosphofructokinases (PFK) are regulatory glycolytic enzymes that convert fructose 6-phosphate and ATP into fructose 1,6-bisphosphate (through PFK-1), fructose 2,6-bisphosphate (through PFK-2), and ADP. Human PFK-1 is tetrameric and isoenzymes include, PFK-1 muscle (PFKM, PFK-A), PFK-1 liver (PFKL, PFK-B), and PFK-1 platelet (PFKP, PFK-C, PFKF). PFK-1 is inhibited by ATP and citrate (from the tricarboxylic acid cycle). PFK-1 undergoes activation in the presence of elevated AMP. The most potent activator is fructose-2,6-bisphosphate, which is produced by PFK-2 from the same substrate, fructose 6-phosphate. PFK-2 is bifunctional and a key regulator for PFK-1. PFK-2 catalyzes the synthesis of fructose-2,6-bisphosphate, and contains fructose-2,6-bisphosphatase activity that catalyzes the degradation of fructose-2,6-bisphosphate. PFK-2 is dimeric and isoenzymes include PFK-2 liver (PFKFB1, PFRX), PFK-2 cardiac (PFKFB2), PFK-2 placental (PFKFB3, inducible PFK-2) and PFK-2 testis (PFKFB4).

**Synonyms:**


F6PK, PFRX, PFK/FBPase 1, PFK-2 liver

**Protein Families:**


Druggable Genome

**Protein Pathways:**

Fructose and mannose metabolism

**Product images:**


Western blot (WB) analysis of [AP20411PU-N]  
PFK-2 liv antibody in extracts from HUVEC cells.



Immunohistochemistry: [AP20411PU-N] PFK-2 liv antibody staining of Paraffin-Embedded Human liver carcinoma tissue.