

Product datasheet for AP20426PU-N

PER3 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: Western blot: 1/500-1/1000.

Immunohistochemistry on Paraffin Sections: 1/50-1/200.

Reactivity: Human

Host: Rabbit

Clonality: Polyclonal

Specificity: This antibody detects endogenous levels of Per3 protein.

(region surrounding Pro2)

Formulation: Phosphate buffered saline (PBS), pH~7.2

State: Aff - Purified

State: Liquid purified Ig fraction (> 95% pure by SDS-PAGE).

Preservative: 0.05% Sodium Azide

Concentration: 1.0 mg/ml

Purification: Affinity Chromatography using epitope-specific immunogen.

Conjugation: Unconjugated

Storage: Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: ~43 kDa

Gene Name: period circadian clock 3

Database Link: Entrez Gene 8863 Human

P56645



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

PER3 Rabbit Polyclonal Antibody - AP20426PU-N

Background:


Biological timepieces called circadian clocks are responsible for the regulation of hormonal rhythms, sleep cycles and other behaviors. The superchiasmatic nucleus (SCN), which is located in the brain, was the first mammalian circadian clock to be discovered. A number of transcription factors appearing to be molecular components of the SCN clock have been identified. Mutations within the Clock gene increase the length of the endogenous period and cause a loss of rhythmicity of circadian oscillations. Three mammalian period proteins, designated Per1, Per2 and Per3, exhibit circadian rhythyms in the SCN. During subjective night, Per1 and Per2 RNA levels increase in response to light pulses while Per3 RNA levels show no change in response to light pulses. Tim, for timeless, interacts with Per1 as well as Per2; and Tim and Per1 negatively regulate Clock-BMAL1-induced transcription.

Synonyms: Period circadian protein homolog 3, hPER3, Circadian clock protein PERIOD 3, GIG13

Protein Families: Druggable Genome, Transcription Factors

Protein Pathways: Circadian rhythm - mammal

Product images:



Western blot analysis of Per3 antibody (Cat.-No.: AP20426PU-N) in extracts from Jurkat cells treated with insulin 0.01U.