

# Product datasheet for AP20443PU-M

## **MELK Rabbit Polyclonal Antibody**

### Product data:

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product Type:	Primary Antibodies
Applications:	IF, IHC, WB
Recommended Dilution:	Western blot: 1/500-1/1000. Immunofluorescence: 1/50-1/200. Immunohistochemistry on Paraffin Sections: 1/50-1/200.
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Specificity:	This antibody detects endogenous levels of MELK protein. (region surrounding asn449)
Formulation:	PBS with 0.02% sodium azide, 50% glycerol, pH7.2 State: Aff - Purified State: Liquid purified Ig fraction (> 95% pure by SDS-PAGE)
Concentration:	1.0 mg/ml
Purification:	Affinity Chromatography using epitope-specific immunogen.
Conjugation:	Unconjugated
Storage:	Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	~75 kDa
Gene Name:	maternal embryonic leucine zipper kinase
Database Link:	<u>Entrez Gene 9833 Human</u> <u>Q14680</u>




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

#### **MELK Rabbit Polyclonal Antibody – AP20443PU-M**


Background: Maternal Embryonic Leucine Zipper Kinase (KIAA0175, HPK38) or MELK, a new member of the Snf1/AMPK family of kinases, encodes a protein with a kinase catalytic domain and a leucine zipper motif consisting of a periodic repetition of leucine residues at every seventh residue located within the N-terminal catalytic domain. This motif has been observed in myriad DNAbinding proteins and is presumed to be involved in protein-DNA interactions, and potentially protein-protein interactions. Research predicts that the gene product of MELK plays a role in the signal transduction events in the egg and early embryo. Mouse and human MELK proteins share 95% sequence identity in the kinase domain and northern blot analysis in mouse indicates that MELK expression is restricted to spermatogonia in the testis and to oocytes in the ovary.

Synonyms:	Maternal embryonic leucine zipper kinase, KIAA0175, hMELK
Protein Families:	Druggable Genome, Protein Kinase

#### **Product images:**



Western blot (WB) analysis of MELK antibody (Cat.-No.: [AP20443PU-N]) in extracts from k562 cells.



Immunohistochemistry (IHC) analyzes of MELK antibody (Cat.-No.: [AP20443PU-N]) in paraffinembedded human liver carcinoma tissue.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US