

Product datasheet for AP20457PU-M

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com

OriGene Technologies, Inc.

EU: info-de@origene.com CN: techsupport@origene.cn

Nucleophosmin (NPM1) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: Western blot: 1/500 - 1/1000.

Immunohistochemistry on paraffin sections: 1/50 - 1/200.

Reactivity: Human, Mouse, Rat

Host: Rabbit

Clonality: Polyclonal

Specificity: This antibody detects endogenous levels of Nucleophosmin protein.

(region surrounding Glu22)

Formulation: Phosphate buffered saline (PBS) with 15 mM sodium azide, approx. pH 7.2.

State: Aff - Purified

State: Liquid purified Ig fraction

Concentration: 1.0 mg/ml

Purification: Affinity chromatography (> 95% (by SDS-PAGE)

Conjugation: Unconjugated

Storage: Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: ~36 kDa

Gene Name: nucleophosmin (nucleolar phosphoprotein B23, numatrin)

Database Link: Entrez Gene 18148 MouseEntrez Gene 25498 RatEntrez Gene 4869 Human

P06748




Background:

The transport of proteins across the nuclear envelope is a selective, multi-step process involving several cytoplasmic factors. Proteins must be recognized as import substrates, dock at the nuclear pore complex and translocate across the nuclear envelope in an ATP-dependent fashion. Several cytosolic and nuclear proteins that are central to this process have been identified. For example, two cytosolic factors critically involved in the recognition and docking process are the karyopherin alpha and karyopherin beta proteins. The karyopherin holoenzyme is a heterodimer of alpha and beta subunits. The nuclear protein B23 (also referred to as nucleophosmin) is involved in ribosomal assembly and rRNA transport. B23 is an abundant protein that is highly phosphorylated by Cdc2 kinase during mitosis.


Synonyms: NPM, NPM1, Numatrin

Protein Families: Druggable Genome, Stem cell - Pluripotency, Transcription Factors

Product images:



Western blot (WB) analysis of Nucleophosmin antibody (Cat.-No.: [AP20457PU-N]) in extracts from Jurkat cells.



Immunohistochemistry (IHC) analyzes of Nucleophosmin antibody (Cat.-No.: [AP20457PU-N]) in paraffin-embedded human breast carcinoma tissue.