

## Product datasheet for AP20463PU-M

## 1 Todact datasticct for Al 204051 0-1

## **DBF4 Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type:** Primary Antibodies

Applications: IHC, WB

Recommended Dilution: Western blot: 1/500 - 1/1000.

Immunohistochemistry on paraffin sections: 1/50 - 1/200.

Reactivity: Human, Mouse

**Host:** Rabbit

Clonality: Polyclonal

**Specificity:** This antibody detects endogenous levels of Dbf4 protein.

(region surrounding Ile18)

**Formulation:** Phosphate buffered saline (PBS) with 0.05% sodium azide, approx. pH 7.2.

State: Aff - Purified

State: Liquid purified Ig fraction

Concentration: 1.0 mg/ml

**Purification:** Affinity chromatography (> 95% (by SDS-PAGE)

Conjugation: Unconjugated

**Storage:** Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

**Stability:** Shelf life: one year from despatch.

**Predicted Protein Size:** ~ 77 kDa

**Gene Name:** DBF4 zinc finger

**Database Link:** Entrez Gene 27214 MouseEntrez Gene 10926 Human

Q9UBU7



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Background:


The Dbf4/Cdc7 protein kinase is essential for the activation of replication origins during S phase. Cdc7-Dbf4 efficiently phosphorylates several proteins that are required for the initiation of DNA replication, including five of the six minichromosome maintenance (Mcm) proteins and the p180 subunit of DNA polymerase alpha-primase. This protein complex consists of the catalytic subunit Cdc7 associating with the regulatory and activating subunit Dbf4, and the kinase activity of the complex is regulated throughout the cell cycle mainly by fluctuating levels of Dbf4. Cdc7 is consistently expressed throughout the cell cycle, while the expression of Dbf4 is absent during G1 phase and accumulates during S and G2 phases. The anaphase-promoting complex rapidly degrades Dbf4 at the time of chromosome segregation, and the stability of Dbf4 remains low during pre-Start G1 phase. The coordinated degradation of Dbf4 and the time of chromosome separation is important to ensuring that prereplicative complexes, which assemble after chromosome segregation, do not immediately refire.

Synonyms: DBF4A, ZDBF1, Chiffon homolog A

**Protein Families:** Druggable Genome

**Protein Pathways:** Cell cycle

## **Product images:**



Western blot (WB) analysis of Dbf4 antibody (Cat.-No.: [AP20463PU-N]) in extracts from NIH-3T3 treated with H2O2 100uM 30'.