

Product datasheet for **AP20486PU-M**

DAP5 (EIF4G2) Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	IF, IHC, WB
Recommended Dilution:	Western blot: 1/500 - 1/1000. Immunohistochemistry on paraffin sections 1/50 - 1/200. Immunofluorescence: 1/50 - 1/200.
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Specificity:	This antibody detects endogenous levels of DAP-5 protein. (region surrounding asn65)
Formulation:	Phosphate buffered saline (PBS) with 0.05% sodium azide, approx. pH 7.2. State: Aff - Purified State: Liquid purified Ig fraction
Concentration:	1.0 mg/ml
Purification:	Affinity chromatography (> 95% (by SDS-PAGE)
Conjugation:	Unconjugated
Storage:	Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	~ 90 kDa
Gene Name:	eukaryotic translation initiation factor 4 gamma 2
Database Link:	<u>Entrez Gene 13690 Mouse</u> <u>Entrez Gene 1982 Human</u> <u>P78344</u>



[View online »](#)

Background:

Death-associated protein 5 (DAP-5) (also known as p97 and NAT1) is a member of the eukaryotic translation initiation factor 4G (eIF4G) family. DAP-5 is ubiquitously expressed and is highly conserved among species. In response to activated FAS or p53, caspase cleaves DAP-5 at position 790 to yield a C-terminal truncated protein which is capable of forming complexes with eIF4A and eIF3. DAP-5 has homology to the carboxy-terminal portion of eIF4G, but lacks the N-terminal region of eIF4G, which is responsible for association with the CAP binding protein eIF4E. By forming translationally inactive complexes with eIF4A and eIF3, but not with eIF4E, DAP-5 functions as a general repressor of translation. During apoptosis, the caspase-activated DAP-5 can mediate CAP-independent translation at least from its own internal ribosome entry site, thus resulting in a positive feedback loop responsible for the continuous translation of DAP-5. DAP-5 is also required for cellular differentiation, as it controls specific gene expression pathways.

Synonyms:


eIF-4-gamma 2, p97, AAG1, NAT1, FLJ41344, Death-associated protein 5

Protein Families:

Transcription Factors

Protein Pathways:

Viral myocarditis

Product images:


Western blot (WB) analysis of DAP-5 antibody (Cat.-No.: [AP20486PU-N]) in extracts from A549 cells.