

## Product datasheet for **AP20551PU-N**

### **DATF1 (DIDO1) Rabbit Polyclonal Antibody**

#### **Product data:**

|                         |   |
|-------------------------|---|
| Product Type:           | Primary Antibodies  |
| Applications:           | IF, IHC, WB   |
| Recommended Dilution:   | <b>Western blot:</b> 1/500-1/1000.<br><b>Immunofluorescence:</b> 1/50-1/200.<br><b>Immunohistochemistry on Paraffin Sections:</b> 1/50-1/200.                       |
| Reactivity:             | Human, Mouse  |
| Host:                   | Rabbit  |
| Clonality:              | Polyclonal  |
| Specificity:            | This antibody detects endogenous levels of DIDO1 / Dio-1 protein.<br>(region surrounding Ser186)  |
| Formulation:            | Phosphate buffered saline (PBS), pH~7.2<br>State: Aff - Purified<br>State: Liquid purified Ig fraction (> 95% pure by SDS-PAGE)<br>Preservative: 0.05% Sodium Azide |
| Concentration:          | 1.0 mg/ml   |
| Purification:           | Affinity Chromatography using epitope-specific immunogen  |
| Conjugation:            | Unconjugated  |
| Storage:                | Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.<br>Avoid repeated freezing and thawing.  |
| Stability:              | Shelf life: one year from despatch.   |
| Predicted Protein Size: | ~244 kDa  |
| Gene Name:              | death inducer-obliterator 1   |
| Database Link:          | <a href="#">Entrez Gene 23856 Mouse</a> <a href="#">Entrez Gene 11083 Human</a><br><a href="#">Q9BTC0</a>   |



[View online »](#)

**Background:**


DIDO1 / Dio-1 (Death Inducer-Obliterator-1) is a putative transcription factor that contains two zinc finger motifs. Dio-1 translocates to the nucleus, and activates apoptosis during limb development. Programmed cell death, a highly regulated form of apoptosis, plays an important role in determining the amount of tissue, the shape, and the definition of each digit during limb development. Dio-1 expression is upregulated when an apoptotic signal is detected, and subsequently apoptosis is induced. This process is similar to the expression of NFκB and NGF in response to external signals. Dio-1 expression is suppressed by caspase inhibitors and Bcl-2 expression. This supports the theory that Dio-1 functions in the onset of programmed cell death.

**Synonyms:**


Death-inducer obliterator 1, DATF1, C20orf158, KIAA0333

**Protein Families:**

Druggable Genome, Transcription Factors

**Product images:**

Western blot (WB) analysis with extracts from 293 cells using DIDO1 / Dio-1 antibody (Cat.-No. AP20551PU-N).



Immunohistochemistry analyzes of paraffin-embedded human colon carcinoma tissue using DIDO1 antibody (Cat.-No. AP20551PU-N).