

# Product datasheet for AP20593PU-M

#### 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

OriGene Technologies, Inc.

### Peroxiredoxin 3 (PRDX3) Rabbit Polyclonal Antibody

**Product data:** 

**Product Type:** Primary Antibodies

Applications: IHC, WB

Recommended Dilution: Western blot: 1/500 - 1/1000.

Immunohistochemistry on paraffin sections: 1/50 - 1/200.

Reactivity: Human, Mouse, Rat

Host: Rabbit
Clonality: Polyclonal

**Specificity:** This antibody detects endogenous levels of PRX III protein.

(region surrounding Lys93)

**Formulation:** Phosphate buffered saline (PBS), pH 7.2.

State: Aff - Purified

State: Liquid purified Ig fraction Preservative: 0.05% sodium azide

**Concentration:** 1.0 mg/ml

**Purification:** Affinity chromatography (> 95% (by SDS-PAGE)

Conjugation: Unconjugated

Storage: Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

**Stability:** Shelf life: one year from despatch.

Predicted Protein Size: ~ 22 kDa

**Gene Name:** peroxiredoxin 3

Database Link: Entrez Gene 11757 MouseEntrez Gene 64371 RatEntrez Gene 10935 Human

P30048






#### Background:

The peroxiredoxin (PRX) family comprises six antioxidant proteins, PRX I, II, III, IV, V and VI, which protect cells from reactive oxygen species (ROS) by preventing the metalcatalyzed oxidation of enzymes. The PRX proteins primarily utilize thioredoxin as the electron donor for antioxidation, although they are fairly promiscuous with regard to the hydroperoxide substrate. In addition to protection from ROS, peroxiredoxins are also involved in cell proliferation, differentiation and gene expression. PRX I, II, IV and VI show diffuse cytoplasmic localization, while PRX III and V exhibit distinct mitochondrial localization. The human PRX I gene encodes a protein that is expressed in several tissues, including liver, kidney, testis, lung and nervous system. PRX II is expressed in testis, while PRX III shows expression in lung. PRX I, II and III are overexpressed in breast cancer and may be involved in its development or progression. Upregulated protein levels of PRX I and II in Alzheimer's disease (AD) and Down syndrome (DS) indicate the involvement of PRX I and II in their pathogenesis.


Synonyms: PRDX-3, Prx3, Prx-3, PRX-III, AOP-1, AOP1, HBC189

**Protein Families:** Transcription Factors

## **Product images:**



Western blot (WB) analysis of PRX III antibody (Cat.-No.: [AP20593PU-N]) in extracts from HeLa cells.



Immunohistochemistry (IHC) analyzes of PRX III antibody (Cat.-No.: [AP20593PU-N]) in paraffinembedded human lung carcinoma tissue.