

## Product datasheet for AP20605PU-S

## **ABCG2 Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type: Primary Antibodies** 

**Applications:** IHC, WB

Recommended Dilution: Western blot: 1/500 - 1/1000.

Immunohistochemistry on paraffin sections: 1/50 - 1/200.

Reactivity: Human Host: Rabbit Clonality:

Polyclonal

Specificity: This antibody detects endogenous levels of ABCG2 protein.

(region surrounding asn338)

Formulation: Phosphate buffered saline (PBS), pH 7.2.

State: Aff - Purified

State: Liquid purified Ig fraction Preservative: 0.05% sodium azide

Concentration: 1.0 mg/ml

Affinity chromatography (> 95% (by SDS-PAGE) **Purification:** 

Conjugation: Unconjugated

Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Storage:

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

**Predicted Protein Size:** ~ 72 kDa

ATP binding cassette subfamily G member 2 (Junior blood group) Gene Name:

Database Link: Entrez Gene 9429 Human

Q9UNQ0



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Background:


ATP-binding cassette (ABC) transporters are an evolutionarily conserved family of proteins that catalyze the transport of molecules across extracellular and intracellular membranes through the energy of ATP hydrolysis. The ABC halftransporter, ABCG2, is also known as placenta-specific ABC transporter and breast cancer resistance protein (BCRP1). ABCG2 confers resistance for a variety of chemotherapeutic agents, including anthracyclines, mitoxantrone, bisantrene and topotecan. Under normal conditions, ABCG2 may serve a protective function by removing toxins from the cell, and plays an important role in regulating stem cell differentiation. ABCG2 is responsible for the side population (SP) phenotype and is widely expressed in a large variety of stem cells, making it an important stem cell marker. ABCG2 may have N-linked glycosylation and may dimerize in vivo. ABCG2 is abundantly expressed in placenta, liver, intestine and stem cells.

**Synonyms:** Breast cancer resistance protein 1, ABCP, MXR


Protein Families: Druggable Genome, ES Cell Differentiation/IPS, Transmembrane

**Protein Pathways:** ABC transporters

## **Product images:**



Western blot (WB) analysis of ABCG2 antibody (Cat.-No.: [AP20605PU-N]) in extracts from HT-29 cells.



Immunohistochemistry (IHC) analyzes of ABCG2 antibody (Cat.-No.: [AP20605PU-N]) in paraffinembedded human brain tissue.