

# Product datasheet for AP20617PU-M

# Leptin (LEP) Rabbit Polyclonal Antibody

### **Product data:**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn


| Product Type:         | Primary Antibodies   |
|-----------------------|--|
| Applications:         | IHC  |
| Recommended Dilution: | Immunohistochemistry on paraffin sections 1/50-1/200.  |
| Reactivity:           | Human, Mouse, Rat  |
| Host:                 | Rabbit   |
| Clonality:            | Polyclonal   |
| Specificity:          | This detects endogenous levels of Ob protein.<br>(region surrounding Lys36)  |
| Formulation:          | Phosphate buffered saline (PBS), pH 7.2<br>State: Aff - Purified<br>State: Liquid purified Ig fraction<br>Preservative: 0.05% sodium azide |
| Concentration:        | 1.0 mg/ml  |
| Purification:         | Affinity-chromatography using epitope-specific immunogen; purity is > 95% (by SDS-PAGE)  |
| Conjugation:          | Unconjugated   |
| Storage:              | Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.<br>Avoid repeated freezing and thawing.                       |
| Stability:            | Shelf life: one year from despatch.  |
| Gene Name:            | leptin   |
| Database Link:        | <u>Entrez Gene 16846 MouseEntrez Gene 25608 RatEntrez Gene 3952 Human</u><br><u>P41159</u>   |




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

|                   | Leptin (LEP) Rabbit Polyclonal Antibody – AP20617PU-M   |
|-------------------|---|
| Background:       | OB Cadherin is a type II classical cadherin from the cadherin superfamily, integral membrane proteins that mediate calcium-dependent cell-cell adhesion. Cadherins are calcium dependent cell adhesion proteins. They preferentially interact with themselves in a homophilic manner in connecting cells; cadherins may thus contribute to the sorting of heterogeneous cell types. Mature cadherin proteins are composed of a large N-terminal extracellular domain, a single membrane-spanning domain, and a small, highly conserved C-terminal cytoplasmic domain. Type II (atypical) cadherins are defined based on their lack of a HAV cell adhesion recognition sequence specific to type I cadherins. Expression of this particular cadherin in osteoblastic cell lines, and its upregulation during differentiation, suggests a specific function in bone development and maintenance. OB Cadherin is expressed in osteoblasts and the brain as well as other tissues.Cadherins constitute a family of transmembrane glycoproteins involved in Ca2+-dependent cell-cell interactions. The members of this family are differentially expressed in various tissues. |
| Synonyms:         | LEP, OB, OBS, Obesity factor, Obese protein   |
| Protein Families: | Druggable Genome, Secreted Protein  |
| Protein Pathway   | <b>s:</b> Adipocytokine signaling pathway, Cytokine-cytokine receptor interaction, Jak-STAT signaling pathway, Neuroactive ligand-receptor interaction  |

## **Product images:**



Immunohistochemistry analyzes of Ob antibody (Cat.-No.: [AP20617PU-N]) in paraffin-embedded human breast carcinoma tissue.



Immunohistochemistry (IHC) analyzes of Ob antibody (Cat.-No.: [AP20617PU-N]) in paraffinembedded human breast carcinoma tissue.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US