

Product datasheet for AP20642PU-M

CD57 / HNK1 Rabbit Polyclonal Antibody

Product data:

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product Type:	Primary Antibodies
Applications:	IHC, WB
Recommended Dilution:	Immunohistochemistry on Paraffin Sections: 1/50-1/200. Western Blot: 1/500-1/1000.
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	Synthetic peptide, corresponding to amino acids 3-50 of Human CD57.
Specificity:	This antibody detects endogenous levels of CD57 protein. (region surrounding Val35)
Formulation:	Phosphate buffered saline (PBS), pH~7.2, 50% Glycerol State: Aff - Purified State: Liquid purified Ig fraction (> 95% pure by SDS-PAGE) Preservative: 0.02% Sodium Azide
Concentration:	1.0 mg/ml
Purification:	Affinity Chromatography using epitope-specific immunogen.
Conjugation:	Unconjugated
Storage:	Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	~38 kDa.



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

CD57 / HNK1 Rabbit Polyclonal Antibody - AP20642PU-MBackground:Over 100 cell surface markers have been identified through the use of monoclonal
antibodies. Many of these markers have proven useful in identifying a specific subpopulation
of cells within a mixed colony. Accordingly, these molecules have been assigned a "cluster of
differentiation" (CD) designation. T lymphocytes displaying the natural killer (NK) cell marker
CD57 (also designated Leu7) on their cell surface are distinguishable from other T cell subsets
by their granular lymphocyte morphology and their clonal expansion in patients with AIDS
and in recipients of bone marrow transplantation. CD57+ cells have also been shown to
localize to sites of certain tumors and large numbers of these cells are detected in the
synovial fluid from patients suffering from rheumatoid arthritis.

Synonyms:

HNK-1, LEU7, NK-1, NK-Cell marker, NK1

Product images:



Immunohistochemistry analyzes of CD57 antibody (Cat.-No.: [AP20642PU-N]) in paraffinembedded human brain tissue.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US