

Product datasheet for **AP20649PU-N**

Livin (BIRC7) Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	WB
Recommended Dilution:	Western blot: 1/500 - 1/1000.
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Specificity:	This antibody detects endogenous levels of ML-IAP protein.
Formulation:	Phosphate buffered saline (PBS), pH 7.2 State: Aff - Purified State: Liquid purified Ig fraction Preservative: 0.05% sodium azide
Concentration:	1.0 mg/ml
Purification:	Affinity-chromatography using epitope-specific immunogen; purity is > 95% (by SDS-PAGE)
Conjugation:	Unconjugated
Storage:	Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	~ 21 kDa
Gene Name:	baculoviral IAP repeat containing 7
Database Link:	Entrez Gene 79444 Human Q96CA5



[View online »](#)

Background:


Inhibitor of apoptosis proteins (IAPs) contain conserved, unique aminoterminal baculovirus IAP repeats (BIRs) and usually a C-terminal RING finger domain. Immunoprecipitation and Western blot analysis indicate that ML-IAP, also known as melanoma inhibitor of apoptosis protein, kidney inhibitor of apoptosis protein (KIAP), livin or BIRC7, binds to caspase-3, -7, and -9, but only inhibits caspase-9. Additionally, ML-IAP physically interacts with SMAC through its BIR domain with a very high affinity and this interaction is very specific. The gene which encodes ML-IAP maps to human chromosome 20q13.3. There is controversy regarding the localization of this protein and its involvement in apoptosis, but it has been suggested that ML-IAP may play a complex role in the regulation of apoptosis.

Synonyms:

KIAP, MLIAP, RNF50

Protein Families:

Druggable Genome

Product images:

Western blot analysis of ML-IAP antibody (Cat.-No.: AP20649PU-N) in extracts from COLO cells.