

## **Product datasheet for AP20704PU-N**

## NF-kB p65 (RELA) Rabbit Polyclonal Antibody

**Product data:** 

**Product Type:** Primary Antibodies

**Applications:** IHC, WB

Recommended Dilution: Western blot: 1/500 - 1/1000.

Immunohistochemistry on paraffin sections: 1/50 - 1/200.

Reactivity: Human, Mouse

Host: Rabbit
Clonality: Polyclonal

Specificity: This antibody detects endogenous levels of NFkB-p65 protein.

(region surrounding Leu523)

**Formulation:** Phosphate buffered saline (PBS), pH 7.2

State: Aff - Purified

State: Liquid purified Ig fraction Preservative: 15 mM sodium azide

**Concentration:** 1.0 mg/ml

**Purification:** Affinity-chromatography using epitope-specific immunogen; purity is > 95% (by SDS-PAGE)

Conjugation: Unconjugated

Storage: Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

**Stability:** Shelf life: one year from despatch.

**Predicted Protein Size:** ~ 60, 75 kDa

**Gene Name:** RELA proto-oncogene, NF-kB subunit

Database Link: Entrez Gene 19697 MouseEntrez Gene 5970 Human

Q04206



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



**Background:** p65 is a subunit of the nuclear factor kappa B. The transcription factor NFkB is widely

recognized as a critical mediator of immune and inflammatory responses. In most cell types, NFkB is found in the cytoplasm where it is associated with an inhibitory prd in many tissues. A high proportion of spontaneous NIH/3T3 transformants overexpress c-Met and by transfection analysis the c-Met proto-oncogene has been shown to exhibit transforming

activity.

Synonyms: NF kappa B p65, NFkB p65, Transcription factor p65, Rel A, NFKB3


**Protein Families:** Druggable Genome, Transcription Factors

**Protein Pathways:** Acute myeloid leukemia, Adipocytokine signaling pathway, Apoptosis, B cell receptor signaling

pathway, Chemokine signaling pathway, Chronic myeloid leukemia, Cytosolic DNA-sensing pathway, Epithelial cell signaling in Helicobacter pylori infection, MAPK signaling pathway, Neurotrophin signaling pathway, NOD-like receptor signaling pathway, Pancreatic cancer, Pathways in cancer, Prostate cancer, RIG-I-like receptor signaling pathway, Small cell lung


cancer, T cell receptor signaling pathway, Toll-like receptor signaling pathway

## **Product images:**



Hela whole cell lysate NFκB-p65 (L523) pAb at 1:500 dilution Western blot (WB) analyzes of NF?B-p65 antibody (Cat.-No.: AP20704PU-N) in extracts from hela cells.





Immunohistochemistry (IHC) analyzes of NF?B-p65 antibody (Cat.-No.: AP20704PU-N) in paraffinembedded human breast carcinoma tissue.