

# Product datasheet for AP20708PU-M

## Product datasneet for AP20/06PU-W

# **EGFR Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type:** Primary Antibodies

Applications: WE

**Recommended Dilution:** Western blot: 1/500 - 1/1000.

Reactivity: Human, Mouse, Rat

**Host:** Rabbit

Clonality: Polyclonal

**Specificity:** This antibody detects endogenous levels of EGFR protein.

(region surrounding Thr1191)

**Formulation:** Phosphate buffered saline (PBS), pH 7.2

State: Aff - Purified

State: Liquid purified lg fraction Preservative: 0.05% sodium azide

Concentration: 1.0 mg/ml

**Purification:** Affinity-chromatography using epitope-specific immunogen; purity is > 95% (by SDS-PAGE)

Conjugation: Unconjugated

**Storage:** Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

**Stability:** Shelf life: one year from despatch.

**Predicted Protein Size:** ~ 175 kDa

**Gene Name:** epidermal growth factor receptor

Database Link: Entrez Gene 13649 MouseEntrez Gene 24329 RatEntrez Gene 1956 Human

P00533



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

#### EGFR Rabbit Polyclonal Antibody - AP20708PU-M

**Background:** The EGF receptor family comprises several related receptor tyrosine kinases that are

frequently overexpressed in a variety of carcinomas. Members of this receptor family include EGFR (HER1), Neu (ErbB-2, HER2), ErbB-3 (HER3), and ErbB-4 (HER4), which form either homodimers or heterodimers upon ligand binding. Exons in the EGFR gene product are frequently either deleted or duplicated to produce deletion mutants (DM) or tandem duplication mutants (TDM), respectively, which are detected at various molecular weights. EGFR binds several ligands, including epidermal growth factor (EGF), transforming growth factor α (TGFα), amphiregulin and heparin binding-EGF (HB-EGF). Ligand binding promotes the internalization of EGFR via Clathrin-coated pits and its subsequent degradation in

response to its intrinsic tyrosine kinase.

Synonyms: Epidermal growth factor receptor, EGF Receptor, erbB-1, c-ErbB-1

Protein Families: Adult stem cells, Cancer stem cells, Druggable Genome, ES Cell Differentiation/IPS, Protein

Kinase, Secreted Protein, Stem cell relevant signaling - JAK/STAT signaling pathway,

Transmembrane

**Protein Pathways:** Adherens junction, Bladder cancer, Calcium signaling pathway, Colorectal cancer, Cytokine-

cytokine receptor interaction, Dorso-ventral axis formation, Endocytosis, Endometrial cancer,

Epithelial cell signaling in Helicobacter pylori infection, ErbB signaling pathway, Focal adhesion, Gap junction, Glioma, GnRH signaling pathway, MAPK signaling pathway, Melanoma, Non-small cell lung cancer, Pancreatic cancer, Pathways in cancer, Prostate

cancer, Regulation of actin cytoskeleton

### **Product images:**



Western blot analyzes of EGFR antibody (Cat.-No.: [AP20708PU-N]) in extracts from HeLa cells.