

Product datasheet for AP20714PU-M

EIF4EBP1 Rabbit Polyclonal Antibody

Product data:

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

| Product Type: | Primary Antibodies | |
|-------------------------|--|--|
| Applications: | IHC, WB | |
| Recommended Dilution: | Western blot: 1/500 - 1/1000. Immunohistochemistry on paraffin sections: 1/50 - 1/200. | |
| Reactivity: | Human, Mouse, Rat | |
| Host: | Rabbit | |
| Clonality: | Polyclonal | |
| Specificity: | This antibody detects endogenous levels of 4E-BP1 protein. (region surrounding Gly39) | |
| Formulation: | Phosphate buffered saline (PBS), pH 7.2 State: Aff - Purified State: Liquid purified lg fraction Preservative: 0.05% sodium azide | |
| Concentration: | 1.0 mg/ml | |
| Purification: | Affinity-chromatography using epitope-specific immunogen; purity is > 95% (by SDS-PAGE) | |
| Conjugation: | Unconjugated | |
| Storage: | Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing. | |
| Stability: | Shelf life: one year from despatch. | |
| Predicted Protein Size: | ~ 15 to 20 kDa | |
| Gene Name: | eukaryotic translation initiation factor 4E binding protein 1 | |
| Database Link: | <u>Entrez Gene 13685 MouseEntrez Gene 116636 RatEntrez Gene 1978 Human</u> <u>Q13541</u> | |



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US


| | EIF4EBP1 Rabbit Polyclonal Antibody – AP20714PU-M |
|-----------------|---|
| Background: | The translation of proteins from eukaryotic mRNA is initiated by the multisubunit complex eIF-4F, which associates with the mRNA 5' cap structure. eIF-4E, a component of eIF-4F, is responsible for binding to the 5' cap structure and for the assembly of the eIF-4F complex. The regulatory protein 4E-BP1, also referred to as PHAS-I, inhibits eIF-4E function. Phosphorylation of 4E-BP1 by S6 kinase p70, MAP kinases or PKCs causes the disassociation of 4E-BP1 from eIF-4E, promoting translation. A protein that is functionally related to 4E-BP1, designated 4E-BP2, also associates with eIF-4E. |
| Synonyms: | PHAS-I, PHAS-1, PHAS1 |
| Protein Pathway | s: Acute myeloid leukemia, ErbB signaling pathway, Insulin signaling pathway, mTOR signaling pathway |

Product images:

~ \$%~

| | -117 -85 |
|-----------|-------------|
| | -49 |
| | -34 |
| 4E-BP1- 🗕 | -25 |

Western blot analyzes of 4E-BP1 antibody (Cat.-No.: [AP20714PU-N]) in extracts from HeLa cells.



Immunohistochemistry analyzes of 4E-BP1 antibody (Cat.-No.: [AP20714PU-N]) in paraffinembedded human breast carcinoma tissue.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US