

## Product datasheet for AP20762PU-N

## 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

CN: techsupport@origene.cn

OriGene Technologies, Inc.

## **TrkB (NTRK2) Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type:** Primary Antibodies

Applications: IHC

Recommended Dilution: Immunohistochemistry on paraffin sections 1/50 - 1/200.

Reactivity: Human, Mouse, Rat

Host: Rabbit
Clonality: Polyclonal

**Immunogen:** Synthetic peptide, corresponding to amino acids 102-158 of Human Sarcoglycan-β.

**Specificity:** This antibody detects endogenous levels of Trk B protein.

(region surrounding Asp699)

**Formulation:** Phosphate buffered saline (PBS), pH 7.2.

State: Aff - Purified

State: Liquid purified Ig fraction Preservative: 0.05% sodium azide

Concentration: 1.0 mg/ml

**Purification:** Affinity chromatography (> 95% (by SDS-PAGE)

Conjugation: Unconjugated

Storage: Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

**Stability:** Shelf life: one year from despatch.

Predicted Protein Size: ~ 92, 145 kDa

**Gene Name:** neurotrophic receptor tyrosine kinase 2

Database Link: Entrez Gene 18212 MouseEntrez Gene 25054 RatEntrez Gene 4915 Human

Q16620





Background:


The Trk proto-oncogene encodes a tyrosine protein kinase, Trk A, also designated Trk gp140, that serves as a receptor for certain neurotrophic factors including nerve growth factor (NGF) and neurotrophin-3 (NT-3). Trk B is a tyrosine kinase gene highly related to Trk A. Trk B expression is confined to tissues within the central and peripheral nervous systems. The brain-derived neurotrophic factor (BDNF) and NT-3, but not NGF, can induce rapid phosphorylation on tyrosine of Trk B gp145, one of the receptors encoded by Trk B, although BDNF elicits a response at least two orders of magnitude greater than NT-3. Thus it appears that Trk B gp145 may represent a neurotrophic receptor for BDNF and NT-3. The third member of the Trk family of tyrosine kinases, Trk C, encodes a protein designated Trk C gp145 that is preferentially expressed in brain tissue, is equally related to Trk A and Trk B, and is a functional receptor for NT-3.

Synonyms: TRKB, NTRK2

**Protein Families:** Druggable Genome, Protein Kinase, Transmembrane

**Protein Pathways:** MAPK signaling pathway, Neurotrophin signaling pathway

## **Product images:**



Immunohistochemistry (IHC) analyzes of Trk B antibody (Cat.-No.: AP20762PU-N) in paraffinembedded human brain tissue.