

## Product datasheet for **AP20769PU-M**

### VEGFD Rabbit Polyclonal Antibody

#### Product data:

Product Type:	Primary Antibodies
Applications:	IHC, WB
Recommended Dilution:	<b>Western blot:</b> 1/500 - 1/1000. <b>Immunohistochemistry on paraffin sections</b> 1/50 - 1/200.
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Specificity:	This antibody detects endogenous levels of VEGF-D protein. (region surrounding Ile202)
Formulation:	Phosphate buffered saline (PBS), pH 7.2. State: Aff - Purified State: Liquid purified Ig fraction Preservative: 0.05% sodium azide
Concentration:	1.0 mg/ml
Purification:	Affinity chromatography (> 95% (by SDS-PAGE)
Conjugation:	Unconjugated
Storage:	Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	~ 40 kDa
Gene Name:	c-fos induced growth factor
Database Link:	<u><a href="#">Entrez Gene 2277 Human</a></u> <u><a href="#">O43915</a></u>

[View online »](#)

**Background:**

The onset of angiogenesis is believed to be an early event in tumorigenesis and may facilitate tumor progression and metastasis. Several growth factors with angiogenic activity have been described. These include fibroblast growth factor (FGF), platelet derived growth factor (PDGF) and vascular endothelial growth factor (VEGF). Several forms of VEGF have been identified, including VEGF, VEGF-B, VEGFC and VEGF-D (also designated FIGF). Characteristic of VEGF proteins, the central region of VEGF-D contains eight cysteine residues. These residues are essential for homodimerization. VEGFD may play a role in tumor progression, as it is induced by c-Fos, which is required for conversion of early stage tumors to malignant tumors. It has been observed that overexpression of VEGF-D induces morphological changes in fibroblasts.

**Synonyms:**


FIGF, VEGF-D, FIGF

**Protein Families:**


Druggable Genome, Secreted Protein

**Protein Pathways:**

Bladder cancer, Cytokine-cytokine receptor interaction, Focal adhesion, mTOR signaling pathway, Pancreatic cancer, Pathways in cancer, Renal cell carcinoma

**Product images:**


Western blot (WB) analyzes of VEGF-D antibody (Cat.-No. [AP20769PU-N]) in extracts from COS-7 cells.



Immunohistochemistry (IHC) analyzes of VEGF-D antibody (Cat.-No. [AP20769PU-N]) in paraffin-embedded human lung carcinoma tissue.