

Product datasheet for AP20789PU-N

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

PGA3 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: Western blot: 1/500-1/1000.

Immunohistochemistry on Paraffin Sections: 1/50-1/200.

Reactivity: Human
Host: Rabbit

Clonality: Polyclonal

Immunogen: Synthetic peptide, corresponding to amino acids 281-330 of Human Pepsin A.

Specificity: This antibody detects endogenous levels of Pepsin protein.

(region surrounding Met307)

Formulation: Phosphate buffered saline (PBS), pH 7.2

State: Aff - Purified

State: Liquid purified Ig fraction (> 95% pure by SDS-PAGE)

Preservative: 0.05% Sodium Azide

Concentration: 1.0 mg/ml

Purification: Affinity Chromatography using epitope-specific immunogen

Conjugation: Unconjugated

Storage: Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: ~42 kDa

Gene Name: pepsinogen 3, group I (pepsinogen A)

Database Link: Entrez Gene 643847 HumanEntrez Gene 643834 Human

PODJD8





Background:


Pepsin is one of the main proteolytic enzymes secreted by the gastric mucosa. Pepsin consists of a single polypeptide chain and arises from its precursor, pepsinogen, by removal of a 41 amino acid segment from the N-terminus. Pepsinogen is synthesized in the stomach lining, and hydrochloric acid, also produced by the gastric mucosa, is necessary to convert the inactive enzyme and to maintain the optimum acidity (pH 1-3) for Pepsin function. Pepsin is particularly effective in cleaving peptide bonds involving aromatic amino acids. Pepsin shows extremely broad specificity; although bonds involving phenylalanine and leucine are preferred, many others are also cleaved to some extent. Pepsin A is a member of the subfamily A1 within the Pepsin family and is the predominant endopeptidase in the gastric juice of vertebrates. Pepsin A is inhibited by ovUS-1, a uterine Serpin.

Synonyms: Pepsin A, PGA, pepsinogen 3 group I, pepsinogen 4 group I, pepsinogen 5 group I, Pepsinogen


A, PGA3, PGA4, PGA5

Protein Families: Druggable Genome

Product images:



Western blot analysis of Pepsin A antiboy Cat.-No AP20789PU-N in extracts from K562 cells.



Immunohistochemistry analysis of Pepsin A antibody Cat.-No AP20789PU-N in paraffinembedded human lung carcinoma tissue.