

## Product datasheet for AP20806PU-M

## 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

CN: techsupport@origene.cn

OriGene Technologies, Inc.

## FOXO4 pSer197 Rabbit Polyclonal Antibody

**Product data:** 

**Product Type:** Primary Antibodies

**Applications:** IHC, WB

Recommended Dilution: Western blot: 1/500 - 1/1000.

Immunohistochemistry on paraffin sections: 1/50 - 1/200.

Reactivity: Human, Mouse

**Host:** Rabbit

Clonality: Polyclonal

**Specificity:** This antibody detects endogenous levels of AFX1 protein only when phosphorylated at

Ser197.

**Formulation:** Phosphate buffered saline (PBS), pH 7.2.

State: Aff - Purified

State: Liquid purified Ig fraction Preservative: 0.05% sodium azide

**Concentration:** 1.0 mg/ml

**Purification:** Affinity chromatography (> 95% (by SDS-PAGE)

Conjugation: Unconjugated

Storage: Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

**Stability:** Shelf life: one year from despatch.

**Predicted Protein Size:** ~ 55, 70 kDa

**Gene Name:** forkhead box O4

Database Link: Entrez Gene 54601 MouseEntrez Gene 4303 Human

P98177






Background:

FKHR (for forkhead in rhabdomyosarcoma), FKHRL1, and AFX1 are members of a subfamily of the forkhead family of transcription factors. AFX1, also known as FoxO4, is expressed in a wide variety of tissues and, like other FKHR proteins, AFX1 contains a single forkhead domain and serineproline-rich region, which mediate DNA binding. AFX1-mediated transcriptional activation is regulated by the serine/threonine kinase Akt1, which phosphorylates AFX1 (in humans, this phosphorylation can occur at several sites, including Ser 197) and in turn, sequesters AFX1 in the cytosol, thereby blocking nuclear localization and DNA binding. Genetic mutations in FKHR genes, including the t(2;13) and t(1;3) translocations, are commonly found in alveolar rhabdomyosarcomas. Additionally, the t(x;11) translocation of the AFX1 gene, which involves the fusion of a serine-prolinerich sequence of AFX1 to the carboxyterminus of a truncated MLL, results in acute lymphocytic leukemia.

Synonyms: Forkhead box protein O4

Protein Families: Transcription Factors

## **Product images:**



Western blot (WB) analyzes of p-AFX1 antibody (Cat.-No.: [AP20806PU-N]) in extracts from COLO205 cells.



Immunohistochemistry (IHC) analyzes of p-AFX1 (Cat.-No.: [AP20806PU-N]) in paraffin-embedded human brain tissue.