

Product datasheet for **AP20815PU-M**

VASP pSer238 Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	IHC, WB
Recommended Dilution:	Western blot: 1/500 - 1/1000. Immunohistochemistry on paraffin sections 1/50 - 1/200.
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Specificity:	This antibody detects endogenous levels of p-VASP protein, only when phosphorylated at Ser238.
Formulation:	Phosphate buffered saline (PBS), pH 7.2 State: Aff - Purified State: Liquid purified Ig fraction Preservative: 0.05% sodium azide
Concentration:	1.0 mg/ml
Purification:	Affinity chromatography (> 95% (by SDS-PAGE)
Conjugation:	Unconjugated
Storage:	Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	~ 48 kDa
Gene Name:	vasodilator-stimulated phosphoprotein
Database Link:	<u>Entrez Gene 22323 Mouse</u> <u>Entrez Gene 361517 Rat</u> <u>Entrez Gene 7408 Human</u> <u>P50552</u>



[View online »](#)

Background:

The Wiskott-Aldrich syndrome (WAS) is characterized by thrombocytopenia, eczema, defects in cell-mediated and humoral immunity and a propensity for lymphoproliferative diseases. The syndrome is the result of a mutation in the gene encoding a proline-rich protein termed WASP. A distantly related protein, VASP (vasodilator-stimulated phosphoprotein), is involved in the maintenance of cytoarchitecture by interacting with Actin-like filaments. VASP shares a limited degree of homology with the amino-terminus of WASP, which is frequently mutated in WAS patients. An established substrate of cAMP and cGMP dependent kinases, VASP is phosphorylated on a regulatory serine residue 157 and localizes to focal adhesions, microfilaments and highly active regions of the plasma membrane. VASP is also phosphorylated on Serine 239 by cGMP-dependent protein kinase.

Synonyms:


Vasodilator-stimulated phosphoprotein

Protein Families:


Druggable Genome, Stem cell - Pluripotency

Protein Pathways:

Fc gamma R-mediated phagocytosis, Focal adhesion, Leukocyte transendothelial migration

Product images:


Western blot (WB) analyzes of p-VASP antibody (Cat.-No.: [AP20815PU-N]) in extracts from NIH/3T3 cells.



Immunohistochemistry (IHC) analyzes of p-VASP antibody (Cat.-No.: [AP20815PU-N]) in paraffin-embedded human cervix carcinoma tissue.