

Product datasheet for **AP20816PU-N**

VAV1 pTyr174 Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	WB
Recommended Dilution:	Western blot: 1/500 - 1/1000.
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Specificity:	This antibody detects endogenous levels of Vav1 protein only when phosphorylated at Tyr174.
Formulation:	Phosphate buffered saline (PBS), pH 7.2. State: Aff - Purified State: Liquid purified Ig fraction Preservative: 0.05% sodium azide
Concentration:	1.0 mg/ml
Purification:	Affinity chromatography (> 95% (by SDS-PAGE)
Conjugation:	Unconjugated
Storage:	Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	~ 100 kDa
Gene Name:	vav guanine nucleotide exchange factor 1
Database Link:	Entrez Gene 22324 Mouse Entrez Gene 25156 Rat Entrez Gene 7409 Human P15498



[View online »](#)

Background:

The Vav gene was originally identified on the basis of its oncogenic activation during the course of gene transfer assays. The major translational product of the Vav protooncogene has been identified as a protein containing an array of structural motifs. Contained within its amino terminus are a helix-loop-helix domain and a leucine zipper motif similar to that of Myc family proteins; deletion of this region of p95Vav causes its oncogenic activation. In addition, p95Vav contains an SH2 domain, which could indicate its role as a substrate for tyrosine kinases. Expression of p95Vav is limited exclusively to cells of hematopoietic origin, including those of the erythroid, lymphoid and myeloid lineages. These results suggest that p95Vav may represent a new type of signal transduction molecule involved in the transduction of tyrosine phosphorylation signaling into transcriptional events.

Synonyms:


VAV-1

Protein Families:

Druggable Genome, Transcription Factors

Protein Pathways:

B cell receptor signaling pathway, Chemokine signaling pathway, Fc epsilon RI signaling pathway, Fc gamma R-mediated phagocytosis, Focal adhesion, Leukocyte transendothelial migration, Natural killer cell mediated cytotoxicity, Regulation of actin cytoskeleton, T cell receptor signaling pathway

Product images:


Western blot (WB) analyzes of p-Vav1 antibody (Cat.-No.: AP20816PU-N) in extracts from Jurkat UV cells.