

Product datasheet for AP20856PU-M

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com

EU: info-de@origene.com

CN: techsupport@origene.cn

OriGene Technologies, Inc.

GJA1 pSer367 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: Western blot: 1/500 - 1/1000.

Immunohistochemistry on paraffin sections: 1/50 - 1/200.

Reactivity: Human, Mouse, Rat

Host: Rabbit
Clonality: Polyclonal

Specificity: This antibody detects endogenous levels of Connexin 43 protein when phosphorylated at

Ser367.

Formulation: Phosphate buffered saline (PBS), pH 7.2.

State: Aff - Purified

State: Liquid purified Ig fraction Preservative: 15 mM sodium azide

Concentration: 1.0 mg/ml

Purification: Affinity chromatography (> 95% (by SDS-PAGE)

Conjugation: Unconjugated

Storage: Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: ~ 43 kDa

Gene Name: gap junction protein alpha 1

Database Link: Entrez Gene 14609 MouseEntrez Gene 24392 RatEntrez Gene 2697 Human

P17302





Background:


The connexins are a group of GAP junction proteins, which form a hexamer to compose a connexon. Clusters of connexons form a GAP junction through which low molecular weight proteins may diffuse from cell to cell. Several mammalian cells with malignant phenotypes exhibit decreased connexin expression and GAP junction communication. There is a decrease in GAP junctional communication in Src transformed cells. The decreased communication appears to be associated with tyrosine phosphorylation of connexin 43. Activated c-Src phosphorylates the C-terminal tail of connexin 43 on residue Tyr 265, resulting in a stable interaction between both proteins leading to inhibition of GAP junctional communication. In addition to tyrosine phosphorylation, connexin 43 has also been shown to be phosphorylated on serine in the absence of Src kinases and on both serine and tyrosine in cells expressing Src kinases such as pp60v-Src and/or c-Src.

Synonyms: Gap junction alpha-1 protein, Connexin-43, GJAL

Protein Families: Druggable Genome, Ion Channels: Other, Transmembrane

Protein Pathways: Arrhythmogenic right ventricular cardiomyopathy (ARVC), Gap junction

Product images:



Jurkat whole cell lysate p-Connexin 43 (S367) pAb at 1:500 dilution Western blot (WB) analyzes of p-Connexin 43 antibody (Cat.-No.: [AP20856PU-N]) in extracts from Jurkat cells.



Immunohistochemistry (IHC) analyzes of p-Connexin 43 antibody (Cat.-No.: [AP20856PU-N]) in paraffin-embedded human brain tissue.