

Product datasheet for AP20859PU-S

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CREB1 pSer129 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

IF, IHC **Applications:**

Recommended Dilution: **Immunohistochemistry on paraffin sections** 1/50 - 1/200.

Immunofluorescence: 1/50 - 1/200.

Reactivity: Human, Mouse, Rat

Host: Rabbit Clonality: Polyclonal

Specificity: This antibody detects endogenous levels of CREB-1 protein only when phosphorylated at

Ser129.

Formulation: Phosphate buffered saline (PBS), pH 7.2.

State: Aff - Purified

State: Liquid purified Ig fraction Preservative: 0.05% sodium azide

Concentration: 1.0 mg/ml

Affinity chromatography (> 95% (by SDS-PAGE) **Purification:**

Conjugation: Unconjugated

Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Storage:

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: ~ 38 kDa

cAMP responsive element binding protein 1 Gene Name:

Entrez Gene 12912 MouseEntrez Gene 81646 RatEntrez Gene 1385 Human Database Link:

P16220



Background:


Eukaryotic gene transcription is regulated by sequencespecific transcription factors that bind modular CIS acting promoter and enhancer elements. The ATF/CREB transcription factor family binds the palindromic cAMP response element (CRE) octanucleotide TGACGTCA. The ATF/CREB family includes CREB-1, CREB-2 (also designated ATF-4), ATF-1, ATF-2 and ATF-3. This family of proteins contain highly divergent N-terminal domains, but share a Cterminal leucine zipper for dimerization and DNA binding. Although CREB can bind to DNA in an unphosphorylated state, it cannot activate transcription. Phosphorylation of CREB on Ser 133 by protein kinase A facilitates its interaction with the CREB-binding protein (CBP) and activates the basal transcription complex. CREB functions in neoglucogenesis through interactions with the nuclear coactivator PGC-1. CREB may play a role in the pathogenesis of type II diabetes and dilated cardiomyopathy.

Synonyms: CREB-1

Protein Families: Druggable Genome, Transcription Factors

Protein Pathways: Antigen processing and presentation, Huntington's disease, Melanogenesis, Prostate cancer

Product images:



Immunohistochemistry (IHC) analyzes of p-CREB-1 antibody (Cat.-No.: [AP20859PU-N]) in paraffinembedded human breast carcinoma tissue.