

Product datasheet for AP20863PU-N

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com

OriGene Technologies, Inc.

EU: info-de@origene.com CN: techsupport@origene.cn

PKR (EIF2AK2) pThr451 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: Western blot: 1/500 - 1/1000.

Immunohistochemistry on paraffin sections: 1/50 - 1/200.

Reactivity: Human

Host: Rabbit

Clonality: Polyclonal

Specificity: This antibody detects endogenous levels of PKR protein only when phosphorylated at Thr451.

Formulation: Phosphate buffered saline (PBS), pH 7.2.

State: Aff - Purified

State: Liquid purified lg fraction Preservative: 0.05% sodium azide

Concentration: 1.0 mg/ml

Purification: Affinity chromatography (> 95% (by SDS-PAGE)

Conjugation: Unconjugated

Storage: Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: Predicted band size: 62 kDa; Observed band size: 68 kDa

Gene Name: eukaryotic translation initiation factor 2 alpha kinase 2

Database Link: Entrez Gene 5610 Human

P19525




Background:

An interferon-inducible, RNA-dependent protein Serine/Threonine kinase, PKR has various designations. Mouse PKR is known as DAI, dsJ, PI kinase, p65, p67 or TIK, whereas human PKR is known as p68 or p69. PKR phosphorylates its substrate, a subunit of protein synthesis initiation factor eIF-2, on Ser 51 to inhibit translation. PKR contains two dsRNA binding motifs required for its activation by dsRNA. Three kinds of regulation of PKR enzymatic activity occur, and these include transcriptional regulation in response to interferon, an autoregulatory mechanism controlling PKR expression at the level of translation, and post-translational regulation by RNA mediated autophosphorylation. Human PKR contains at least 15 autophosphorylation sites, but only Thr 446 and Thr 451 in the activation loop are critical for its kinase activity. Thr 446 is the in vivo autophosphorylation site of PKR.


Synonyms: elF-2A protein kinase 2, PRKR

Protein Families: Druggable Genome, Protein Kinase, Transcription Factors

Product images:



Western blot (WB) analysis of p-PKR antibody (Cat.-No.: AP20863PU-N) at 1/500 dilution Lane 1:Hela whole cell lysate treated with PMA Lane 2:Raw264.7 whole cell lysate treated with H2O2 Lane 3:NIH-3T3 whole cell lysate treated with LPS



Immunohistochemistry (IHC) analyzes of p-PKR antibody (Cat.-No.: AP20863PU-N) in paraffinembedded human colon carcinoma tissue.