

## Product datasheet for **AP20928PU-M**

### CDK1 pTyr15 Rabbit Polyclonal Antibody

#### Product data:


Product Type:	Primary Antibodies
Applications:	IHC
Recommended Dilution:	<b>Immunohistochemistry on paraffin sections</b> 1/50 - 1/200.
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Specificity:	This antibody detects endogenous levels of p-CDC2 protein only when phosphorylated at Tyr15.
Formulation:	Phosphate buffered saline (PBS, pH 7.2. State: Aff - Purified State: Liquid purified Ig fraction Preservative: 0.05% sodium azide
Concentration:	1.0 mg/ml
Purification:	Affinity chromatography (> 95% (by SDS-PAGE)
Conjugation:	Unconjugated
Storage:	Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	~ 34 kDa
Gene Name:	cyclin-dependent kinase 1
Database Link:	<a href="#">Entrez Gene 12534 Mouse</a> <a href="#">Entrez Gene 54237 Rat</a> <a href="#">Entrez Gene 983 Human</a> <a href="#">P06493</a>



[View online »](#)

<b>Background:</b>	Cdc2, an evolutionarily conserved serine/threonine-specific protein kinase, is essential in the cell cycle transition from G2 to M phase. Cdc2 is regulated by association with B-type cyclins and by reversible phosphorylation. Cyclin B binding facilitates the phosphorylation of Cdc2 p34 on three regulatory sites: threonine 14, tyrosine 15, and threonine 161. In higher eukaryotes, Cdc2 is negatively regulated by phosphorylation of two residues located in the ATP-binding site, Thr 14 and Tyr 15. Cdc2 is positively regulated by the cyclin-dependent phosphorylation of Thr 161. Both phosphorylation and de- phosphorylation at Thr 161 are required for progression through the cell cycle.
<b>Synonyms:</b>	CDK1, CDC2, CDC28A, CDKN1, P34CDC2, p34 protein kinase
<b>Protein Families:</b>	Druggable Genome, Protein Kinase, Stem cell - Pluripotency
<b>Protein Pathways:</b>	Cell cycle, Gap junction, Oocyte meiosis, p53 signaling pathway, Progesterone-mediated oocyte maturation

### Product images:



Immunohistochemistry (IHC) analyzes of p-CDC2 (pTyr15) antibody (Cat.-No.: [AP20928PU-NJ]) in paraffin-embedded human breast carcinoma tissue.