

## **Product datasheet for AP20928PU-N**

### 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com

OriGene Technologies, Inc.

#### EU: info-de@origene.com CN: techsupport@origene.cn

# **CDK1 pTyr15 Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type:** Primary Antibodies

Applications: IHC

Recommended Dilution: Immunohistochemistry on paraffin sections 1/50 - 1/200.

Reactivity: Human, Mouse, Rat

Host: Rabbit
Clonality: Polyclonal

**Specificity:** This antibody detects endogenous levels of p-CDC2 protein only when phosphorylated at

Tyr15.

**Formulation:** Phosphate buffered saline (PBS, pH 7.2.

State: Aff - Purified

State: Liquid purified lg fraction Preservative: 0.05% sodium azide

**Concentration:** 1.0 mg/ml

**Purification:** Affinity chromatography (> 95% (by SDS-PAGE)

Conjugation: Unconjugated

**Storage:** Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

**Stability:** Shelf life: one year from despatch.

**Predicted Protein Size:** ~ 34 kDa

**Gene Name:** cyclin-dependent kinase 1

Database Link: Entrez Gene 12534 MouseEntrez Gene 54237 RatEntrez Gene 983 Human

P06493

### CDK1 pTyr15 Rabbit Polyclonal Antibody - AP20928PU-N

Background: Cdc2, an evolutionarily conserved serine/threonine-specific protein kinase, is essential in the

> cell cycle transition from G2 to M phase. Cdc2 is regulated by association with B-type cyclins and by reversible phosophorylation. Cyclin B binding facilitates the phosphorylation of Cdc2 p34 on three regulatory sites: threonine 14, tyrosine 15, and threonine 161. In higher eukaryotes, Cdc2 is negatively regulated by phosphorylation of two residues located in the

ATP-binding site, Thr 14 and Tyr 15. Cdc2 is positively regulated by the cyclin-dependent phosphorylation of Thr 161. Both phosphorylation and de-phosphorylation at Thr 161 are

required for progression through the cell cycle.


Synonyms: CDK1, CDC2, CDC28A, CDKN1, P34CDC2, p34 protein kinase

Druggable Genome, Protein Kinase, Stem cell - Pluripotency **Protein Pathways:** Cell cycle, Gap junction, Oocyte meiosis, p53 signaling pathway, Progesterone-mediated

oocyte maturation

### **Product images:**

**Protein Families:** 



Immunohistochemistry (IHC) analyzes of p-CDC2 (pTyr15) antibody (Cat.-No.: AP20928PU-N) in paraffin-embedded human breast carcinoma tissue.