

## Product datasheet for **AP20934PU-N**

### **p38 (CRK) (CRK-II, pTry221) Rabbit Polyclonal Antibody**

#### **Product data:**

Product Type:	Primary Antibodies
Applications:	IHC
Recommended Dilution:	<b>Immunohistochemistry on paraffin sections</b> 1/50 - 1/200.
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Specificity:	This antibody detects endogenous levels of p-Crk2 protein only when phosphorylated at Try221.
Formulation:	Phosphate buffered saline (PBS), pH 7.2. State: Aff - Purified State: Liquid purified Ig fraction Preservative: 0.05% sodium azide
Concentration:	1.0 mg/ml
Purification:	Affinity chromatography (> 95% (by SDS-PAGE)
Conjugation:	Unconjugated
Storage:	Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	~ 42 kDa
Gene Name:	v-crk avian sarcoma virus CT10 oncogene homolog
Database Link:	<a href="#">Entrez Gene 12928 Mouse</a> <a href="#">Entrez Gene 54245 Rat</a> <a href="#">Entrez Gene 1398 Human P46108</a>



[View online »](#)

**Background:**

The Crk family of adapter proteins including Crk II, Crk I and Crk L consist mostly of SH2 and SH3 domains. Through the interactions between SH2 domain and phosphotyrosine residues and/or between SH3 domain and proline- rich motifs, they are involved in a variety of signaling cascades. Crk I and Crk II are encoded by the same gene, which undergoes alternative splicing to yield these two proteins, but differ in their biological activities. Crk II has less transforming activity than Crk I, although both Crk I and Crk II bind to many tyrosine-phosphorylated proteins that bind to GRB2. In addition, Crk II becomes rapidly tyrosine-phosphorylated in response to stimulation with insulin-like growth factor-I (IGFI) and might be involved in the IGF-I receptor signalling pathway. The gene encoding Crk I and II maps to human chromosome 17p13, a region which demonstrates frequent deletion or loss of heterozygosity in a wide range of human cancers.

**Synonyms:**


Proto-oncogene C-crk, p38

**Protein Families:**

Druggable Genome, Transcription Factors

**Protein Pathways:**

Chemokine signaling pathway, Chronic myeloid leukemia, ErbB signaling pathway, Fc gamma R-mediated phagocytosis, Focal adhesion, Insulin signaling pathway, MAPK signaling pathway, Neurotrophin signaling pathway, Pathways in cancer, Regulation of actin cytoskeleton, Renal cell carcinoma

**Product images:**

Immunohistochemistry (IHC) analyzes of p-Crk2 antibody (Cat.-No.: AP20934PU-N) in paraffin-embedded human lung adenocarcinoma tissue.