

Product datasheet for **AP20959PU-N**

NMDAR1 (GRIN1) pSer897 Rabbit Polyclonal Antibody

Product data:

| | |
|------------------------------|--|
| Product Type: | Primary Antibodies |
| Applications: | IHC |
| Recommended Dilution: | Immunohistochemistry on paraffin sections: 1:50 - 1:200. |
| Reactivity: | Human, Mouse, Rat |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Specificity: | This antibody detects endogenous levels of p-NMDAR1 (pSer897) protein. |
| Formulation: | Phosphate buffered saline (PBS) with 15 mM sodium azide, approx. pH 7.2 State: Aff - Purified State: Liquid Ig fraction |
| Concentration: | 1.0 mg/ml |
| Purification: | Affinity-chromatography using epitope-specific immunogen, purity is > 95% (by SDS-PAGE) |
| Conjugation: | Unconjugated |
| Storage: | Store at 2 - 8 °C for up to one month or (in aliquots) at -20 °C for longer. Avoid repeated freezing and thawing. |
| Stability: | Shelf life: one year from despatch. |
| Gene Name: | glutamate ionotropic receptor NMDA type subunit 1 |
| Database Link: | <u>Entrez Gene 14810 Mouse</u> <u>Entrez Gene 24408 Rat</u> <u>Entrez Gene 2902 Human</u> <u>Q05586</u> |



[View online »](#)

Background:

NMDA receptor subtype of glutamate-gated ion channels possesses high calcium permeability and voltage-dependent sensitivity to magnesium. Mediated by glycine. Plays a key role in synaptic plasticity, synaptogenesis, excitotoxicity, memory acquisition and learning. It mediates neuronal functions in glutamate neurotransmission. Is involved in the cell surface targeting of NMDA receptors. The ion channels activated by glutamate are divided into two classes. Those that are sensitive to N-methyl-D-aspartate (NMDA) are designated NMDA receptors (NMDAR) while those activated by kainate and α-amino-3-hydroxy-5-methyl-4-isoxalone propionic acid (AMPA) are known as kainate/AMPA receptors (K/AMPA). NMDA receptors are among the most studied receptors in neuroscience because they are involved in neuronal cell development and plasticity, a cellular correlate for learning. NMDA receptors are also implicated in several disorders of the central nervous system including epilepsy and ischemic neuronal cell death. NMDA receptors also appear to be a target for ethanol at physiological concentrations and therefore may play a significant role in alcoholism.

Synonyms:


NMDAR1, GRIN1

Protein Families:

Druggable Genome, Ion Channels: Glutamate Receptors, Transmembrane

Protein Pathways:

Alzheimer's disease, Amyotrophic lateral sclerosis (ALS), Calcium signaling pathway, Huntington's disease, Long-term potentiation, Neuroactive ligand-receptor interaction

Product images:


Immunohistochemistry analyzes of p-NMDAR1 antibody (AP20959PU-N) in paraffin-embedded human brain tissue.