

# Product datasheet for AP20965PU-S

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

## **ATM Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type: Primary Antibodies** 

IHC **Applications:** 

Recommended Dilution: Immunohistochemistry on paraffin sections 1/50 - 1/200.

Reactivity: Rabbit Host: Clonality: Polyclonal

Synthetic peptide, corresponding to amino acids 1957-2006 of Human ATM. Immunogen:

Specificity: This antibody detects endogenous levels of ATM protein.

(region surrounding Ile1987)

Formulation: Phosphate buffered saline (PBS), pH 7.2.

State: Aff - Purified

State: Liquid purified Ig fraction Preservative: 0.05% sodium azide

Concentration: 1.0 mg/ml

**Purification:** Affinity chromatography (> 95% (by SDS-PAGE)

Conjugation: Unconjugated

Storage: Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

**Predicted Protein Size:** ~ 350 kDa

Gene Name: ATM serine/threonine kinase

Database Link: Entrez Gene 472 Human

Q13315



#### ATM Rabbit Polyclonal Antibody - AP20965PU-S

Background:


The phosphatidylinositol kinase (PIK) family members fall into two distinct subgroups. The first subgroup contains proteins such as the PI 3- and PI 4- kinases and the second group comprises the PIK-related kinases. The PIKrelated kinases include Atm, DNA-PKCS and FRAP. These proteins have in common a region of homology at their carboxy-termini that is not present in the PI 3- and PI 4-kinases. The Atm gene is mutated in the autosomal recessive disorder ataxia telangiectasia (AT) that is characterized by cerebellar degeneration (ataxia) and the appearance of dilated blood vessels (telangiectases) in the conjunctivae of the eyes. AT cells are hypersensitive to ionizing radiation, impaired in mediating the inhibition of DNA synthesis and display delays in p53 induction.

Synonyms: Serine-protein kinase ATM, Ataxia telangiectasia mutated, A-T mutated, ATDC, TEL1, TEL01

**Protein Families:** Druggable Genome, Protein Kinase, Transcription Factors

**Protein Pathways:** Apoptosis, Cell cycle, p53 signaling pathway

### **Product images:**



Immunohistochemistry (IHC) analyzes of ATM antibody in paraffin-embedded human lung adenocarcinoma tissue.