

# Product datasheet for AP20970PU-M

# CAMK2A Rabbit Polyclonal Antibody

## **Product data:**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product Type:	Primary Antibodies
21	
Applications:	IHC, WB
Recommended Dilution:	Western blot: 1/500 - 1/1000. Immunohistochemistry on paraffin sections: 1/50 - 1/200.
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Specificity:	This antibody detects endogenous levels of CaMK2-a protein. (region surrounding Lys292)
Formulation:	Phosphate buffered saline (PBS), pH 7.2. State: Aff - Purified State: Liquid purified lg fraction Preservative: 0.05% sodium azide
Concentration:	1.0 mg/ml
Purification:	Affinity chromatography (> 95% (by SDS-PAGE)
Conjugation:	Unconjugated
Storage:	Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	~ 54 kDa
Gene Name:	calcium/calmodulin dependent protein kinase II alpha
Database Link:	<u>Entrez Gene 815 Human</u> <u>Q9UQM7</u>




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

#### **GRIGENE** CAMK2A Rabbit Polyclonal Antibody – AP20970PU-M


Background:The multifunctional CAMK2, or Ca2+/calmodulindependent protein kinase II, is a well known<br/>effector of calcium- and calmodulin- mediated functions. It is present in many tissues but is<br/>most abundant in the brain. CAMK2 is composed of four different chains: alpha, beta,<br/>gamma, and delta. The different isoforms assemble into homo- or heteromultimeric<br/>holoenzymes composed of 8 to 12 subunits. Autophosphorylation plays an important role in<br/>the regulation of the kinase activity. CAMK2 is required for synaptic plasticity, as in Long Term<br/>Potentiation (LTP), a cellular model for learning and memory. CAMK2B is the beta chain.<br/>There are seven named isoforms produced by alternative splicing. The variable region of the<br/>CAMK2B protein is encoded by at least 7 exons (V1 to V7). Alternative splicing within this<br/>region gives rise to CAMK2B isoforms.CAMK2G is the gamma chain. CAMK2G has six<br/>alternatively spliced variants that encode six different isoforms. Some of these variants have<br/>been identified in human tumors.

Synonyms: CAMKA, KIAA0968, CaM-kinase II alpha chain, CaM kinase II subunit alpha, CaMK-II subunit alpha

### **Product images:**



Western blot (WB) analyzes of CaMK2-a antibody (Cat.-No.: [AP20970PU-N]) in extracts from A431 cells.



Immunohistochemistry (IHC) analyzes of CaMK2-a antibody (Cat.-No.: [AP20970PU-N]) in paraffinembedded human A431 tissue.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US