

## Product datasheet for **AP21029PU-N**

### FGFR1 Rabbit Polyclonal Antibody

#### Product data:

Product Type:	Primary Antibodies
Applications:	IHC
Recommended Dilution:	<b>Immunohistochemistry on paraffin sections</b> 1/50-1/200.
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Specificity:	This antibody detects endogenous levels of FGFR1 protein. (region surrounding Glu160)
Formulation:	Phosphate buffered saline (PBS), pH~7.2 State: Aff - Purified State: Liquid purified Ig fraction (> 95% pure by SDS-PAGE) Preservative: 0.05% Sodium Azide
Concentration:	1.0 mg/ml
Purification:	Affinity Chromatography using epitope-specific immunogen
Conjugation:	Unconjugated
Storage:	Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	~ 120, 145 kDa
Gene Name:	fibroblast growth factor receptor 1
Database Link:	<a href="#">Entrez Gene 14182 Mouse</a> <a href="#">Entrez Gene 79114 Rat</a> <a href="#">Entrez Gene 2260 Human P11362</a>



[View online »](#)

**Background:**

FGFR1 (fibroblast growth factor receptor 1) is a member of the fibroblast growth factor receptor family and contains an Ig-like domain and a tyrosine kinase domain. This receptor has multiple isoforms and is a Type I membrane protein. FGFR1 is widely expressed, with distinct isoforms expressed in specific tissues. FGFR1 binds fibroblast growth factor and induces mitogenesis and cellular differentiation. Defects in FGFR1 result in Pfeiffer syndrome associated with craniosynostosis. FGFR1 can be modified by phosphorylation and can bind basic/acidic fibroblast factor depending on the receptor isoform. FGFR1 has been shown to interact with Ncadherin and NCAM. Fibroblast growth factors (FGFs) are members of a large family of structurally related polypeptides (17-38 kDa) that are potent physiological regulators of growth and differentiation of a wide variety of cells of mesodermal, ectodermal and endodermal origin. FGFs are substantially involved in normal development, wound healing and repair, angiogenesis, a variety of neurotrophic activities, in hematopoiesis as well as in tissue remodeling and maintenance. They also have been implicated in pathological conditions such as tumorigenesis and metastasis.

**Synonyms:**


BFGFR, CEK, FGFBR, FLG, FLT2, HBGFR, BFGFR, bFGF-R-1, FLT-2, N-sam, Proto-oncogene c-Fgr

**Protein Families:**

Druggable Genome, Protein Kinase, Transmembrane

**Protein Pathways:**

Adherens junction, MAPK signaling pathway, Melanoma, Pathways in cancer, Prostate cancer, Regulation of actin cytoskeleton

**Product images:**

Immunohistochemistry analysis of FGFR1 antibody (Cat.-No.: AP21029PU-N) in paraffin-embedded human lung adenocarcinoma tissue.