

Product datasheet for AP21056PU-M

PAK1 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IF, IHC, WB

Recommended Dilution: Western blot: 1/500 - 1/1000.

Immunohistochemistry on paraffin sections: 1/50 - 1/200.

Immunofluorescence: 1/50 - 1/200.

Reactivity: Human, Mouse, Rat

Host: Rabbit

Clonality: Polyclonal

Specificity: This antibody detects endogenous levels of PAKalpha protein.

(region surrounding Glu195)

Formulation: Phosphate buffered saline (PBS), pH 7.2.

State: Aff - Purified

State: Liquid purified Ig fraction Preservative: 0.05% sodium azide

Concentration: 1.0 mg/ml

Purification: Affinity chromatography (> 95% (by SDS-PAGE)

Conjugation: Unconjugated

Storage: Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: ~ 60 kDa

Gene Name: p21 (RAC1) activated kinase 1

Database Link: Entrez Gene 18479 MouseEntrez Gene 29431 RatEntrez Gene 5058 Human

Q13153



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Background:

Pak1 belongs to the Ser/Thr protein kinase family and may function by modifying and partially stabilizing thermolabile DNA polymerases, perhaps during DNA repair. Expression of PAK1 increases specific activity of p53 in DNA-binding assays accompanied by a corresponding increase in transactivation. In vitro protein kinase assays show that GST-Pak1 can autophosphorylate, and can phosphorylate casein as an exogenous substrate. The Pak1 protein autophosphorylates on serine residues and preferentially binds to activated Cdc42p both in vitro and in vivo. This binding is mediated through the p21 binding domain on Pak1p and the effector domain on Cdc42p. Pak1 catalytic domain binds to the same highly conserved region on the regulatory domain that binds Cdc42, a GTPase protein capable of activating Pak1.

Synonyms:

PAK 1, PAK-1, Alpha-PAK, PAK alpha, p21-activated kinase 1, p65-PAK


Protein Families:

Druggable Genome, Protein Kinase, Stem cell - Pluripotency

Protein Pathways:


Axon guidance, Chemokine signaling pathway, Epithelial cell signaling in Helicobacter pylori infection, ErbB signaling pathway, Fc gamma R-mediated phagocytosis, Focal adhesion, MAPK signaling pathway, Natural killer cell mediated cytotoxicity, Regulation of actin cytoskeleton, Renal cell carcinoma, T cell receptor signaling pathway

Product images:



Western blot (WB) analysis of PAKalpha antibody (Cat.-No.: [AP21056PU-N]) in extracts from K562 cells.





Immunohistochemistry (IHC) analyzes of PAKalpha antibody (Cat.-No.: [AP21056PU-N]) in paraffin-embedded human breast carcinoma tissue.