

## **Product datasheet for AP21101PU-N**

# **CDC25A Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type:** Primary Antibodies

Applications: WE

Recommended Dilution: Western blot: 1/500 - 1/1000.

Reactivity: Human, Mouse, Rat

**Host:** Rabbit

Clonality: Polyclonal

**Specificity:** This antibody detects endogenous levels of Cdc25A protein.

**Formulation:** Phosphate buffered saline (PBS), pH 7.2.

State: Aff - Purified

State: Liquid purified Ig fraction Preservative: 0.05% sodium azide

**Concentration:** 1.0 mg/ml

**Purification:** Affinity chromatography (> 95% (by SDS-PAGE)

**Conjugation:** Unconjugated

Storage: Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

**Stability:** Shelf life: one year from despatch.

Predicted Protein Size: ~ 59 kDa

**Gene Name:** cell division cycle 25A

Database Link: Entrez Gene 12530 MouseEntrez Gene 171102 RatEntrez Gene 993 Human

P30304



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

#### CDC25A Rabbit Polyclonal Antibody - AP21101PU-N

#### Background:


The Cdc2/cyclin B enzyme, involved in regulation of mitosis in eukaryotic cells, is subject to multiple levels of control. Among these, the regulation of the catalytic subunit by tyrosine phosphorylation is the best understood. tyrosine phosphorylation inhibits the Cdc2/cyclin B complex, while tyrosine dephosphorylation, which occurs at the onset of mitosis, directly activates the pre-MPH complex. The Cdc25 gene serves as a rate-limiting mitotic activator, apparently due to its action as the Cdc2 tyrosine phosphatase. In the absence of Cdc25, Cdc2 accumulates in a tyrosine phosphorylated state. In addition, Cdc25 proteins from a variety of species have been shown to share a low degree of sequence similarity with other tyrosine phosphatases. The Cdc25 gene family consists of at least three members that share approximately 40% identity in their most conserved carboxy terminal sequences.

Synonyms: CDC-25A

**Protein Families:** Druggable Genome, Phosphatase

**Protein Pathways:** Cell cycle, Progesterone-mediated oocyte maturation

### **Product images:**



Western blot (WB) analysis of Cdc25A antibody (Cat.-No.: AP21101PU-N) in extracts from Jurkat cells.