

Product datasheet for **AP21104PU-N**

HSF1 Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	IHC, WB
Recommended Dilution:	Western blot: 1/500 - 1/1000. Immunohistochemistry on paraffin sections 1/50 - 1/200.
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Specificity:	This antibody detects endogenous levels of HSF1 protein. (region surrounding Asp135)
Formulation:	Phosphate buffered saline (PBS), pH 7.2. State: Aff - Purified State: Liquid purified Ig fraction Preservative: 0.05% sodium azide
Concentration:	1.0 mg/ml
Purification:	Affinity chromatography (> 95% (by SDS-PAGE)
Conjugation:	Unconjugated
Storage:	Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	~ 57, 82 kDa
Gene Name:	heat shock transcription factor 1
Database Link:	<u>Entrez Gene 15499 Mouse</u> <u>Entrez Gene 3297 Human</u> <u>Q00613</u>



[View online »](#)

Background:


Heat shock transcription factors (HSF, also designated HSTF) 1 and 2 are involved in this regulation. HSF1 and HSF2 are upregulated by estrogen at both the mRNA and protein level. HSF1 is normally found as a monomer whose transcriptional activity is repressed by constitutive phosphorylation. Upon activation, HSF1 forms trimers, gains DNA binding activity and is translocated to the nucleus. HSF2 activity is associated with differentiation and development, and, like HSF1, binds DNA as a trimer. Both HSF1 and HSF2 are known to be induced by proteasome inhibitors of the ubiquitin pathway.

Synonyms:


HSTF1, HSF-1

Protein Families:

Transcription Factors

Product images:


Western blot (WB) analysis of HSF1 antibody (Cat.-No.: AP21104PU-N) in extracts from K562 cells.



Immunohistochemistry (IHC) analyzes of HSF1 antibody (Cat.-No.: AP21104PU-N) in paraffin-embedded human colon carcinoma tissue.