

## **Product datasheet for AP21131PU-M**

## Product data:

**Product Type:** Primary Antibodies

**GRK2 Rabbit Polyclonal Antibody** 

Applications: IHC, WB

Recommended Dilution: Western blot: 1/500 - 1/1000.

Immunohistochemistry on paraffin sections: 1/50 - 1/200.

Reactivity: Human, Mouse, Rat

**Host:** Rabbit

**Clonality:** Polyclonal

**Specificity:** This antibody detects endogenous levels of GRK 2 protein.

(region surrounding Val678)

**Formulation:** Phosphate buffered saline (PBS), pH 7.2.

State: Aff - Purified

State: Liquid purified Ig fraction Preservative: 0.05% sodium azide

**Concentration:** 1.0 mg/ml

**Purification:** Affinity chromatography (> 95% (by SDS-PAGE)

Conjugation: Unconjugated

Storage: Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

**Stability:** Shelf life: one year from despatch.

Predicted Protein Size: ~ 80 kDa

**Gene Name:** G protein-coupled receptor kinase 2

Database Link: Entrez Gene 25238 RatEntrez Gene 110355 MouseEntrez Gene 156 Human

P25098



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Background:


Heterotrimeric G protein-mediated signal transduction is a dynamically regulated process with the intensity of signal decreasing over time despite the continued presence of the agonist. This phenomenon, referred to as agonistmediated desensitization, involves phosphorylation of the receptor by two classes of enzymes. The first are the second messenger-regulated kinases such as c-AMP dependent protein kinase A and protein kinase C. The second are the G protein-coupled receptor kinases (GRKs). At least seven members of the GRK family have been identified. These include rhodopsin kinase, GRK 1; two forms of  $\beta$ -adrenergic receptor kinase, GRK 2 ( $\beta$ ARK,  $\beta$ ARK1) and GRK 3 ( $\beta$ ARK2); IT-11 (GRK 4); GRK 5, GRK 6 and GRK 7. Phosphorylation of receptors by GRKs appears to be strictly dependent on the receptor being in its agonist-activated state.

Synonyms: BARK, BARK1, GRK2, Beta-adrenergic receptor kinase 1


**Protein Families:** Druggable Genome, Protein Kinase

**Protein Pathways:** Chemokine signaling pathway, Endocytosis

## **Product images:**



Western blot (WB) analysis of GRK 2 antibody (Cat.-No.: [AP21131PU-N]) in extracts from Jurkat cells.



Immunohistochemistry (IHC) analyzes of GRK 2 antibody (Cat.-No.: [AP21131PU-N]) in paraffinembedded human colon carcinoma tissue.