

Product datasheet for AP21133PU-N

LOC101930123 Rabbit Polyclonal Antibody

Product data:

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product Type:	Primary Antibodies
Applications:	WB
Recommended Dilution:	Western Blot: 1/500-1/1000.
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Specificity:	This antibody detects endogenous levels of eEF2K protein. (region surrounding Gly358)
Formulation:	Phosphate buffered saline (PBS), pH 7.2. State: Aff - Purified State: Liquid purified lg fraction (> 95% by SDS-PAGE) Preservative: 0.05% Sodium Azide
Concentration:	1.0 mg/ml
Purification:	Affinity Chromatography using epitope-specific immunogen
Conjugation:	Unconjugated
Storage:	Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	~ 105 kDa
Database Link:	<u>Entrez Gene 13631 MouseEntrez Gene 25435 RatEntrez Gene 101930123 Human</u> <u>O00418</u>




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

CRIGENE LOC101930123 Rabbit Polyclonal Antibody – AP21133PU-N

Background:The activity of the purified eukaryotic elongation-factor-2 kinase (eEF2K) is completely
dependent on calcium and calmodulin, and autophosphorylation on serine and threonine
residues is calcium/calmodulin-dependent. eEF2K is a ubiquitous protein kinase that
phosphorylates and inactivates eEF2, and thus can modulate the rate of polypeptide chain
elongation during translation. eEF2K is 120 kDa protein that is detected in skeletal muscle
extracts and is phosphorylated rapidly by SAPK4, but poorly by p38, p38g, JNK or ERK 2.
SAPK4 phosphorylates eEF2K at Ser 359 and Ser 396 in vitro, causing its inactivation. The
phosphorylation of eEF2K at Ser 359 is also induced by insulin-like growth factor-1. Ser 359 is
in close proximity to Ser 366 and the Ser 366 residue also becomes phosphorylated in
response to growth factors. eEF2K is phosphorylated by p70 S6 kinase at Ser 366 and this
results in the inactivation of eEF2K, especially at low (micromolar) calcium concentrations.

Synonyms:	Elongation factor 2 kinase, EEF2K, EF2K
Protein Families:	Druggable Genome, Protein Kinase

Product images:



Western blot (WB) analysis of eEF2K antibody (Cat.-No.: AP21133PU-N) in extracts from Jurkat cells.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US